

## **PRODUCT INFORMATION**

Warning: Undefined variable ShasAttributeValueDescription in C:\www.root\mirror.dimablo.com\wp-content\plugins\woocommerce-print-products\public:class-woocommerce-print-products-public-php on line 2806 bM69 Clone ID

CD244;2B4;SLAMF4;NKR2B4;NAIL;h2B4

Host Species

Rabbit Anti-2B4 antibody(DM69); Rabbit mAb Description Delivery 3~4 weeks

Q9BZW8 Uniprot ID IgG type Rabbit IgG Clonality Monoclonal Reactivity ELISA; Flow Cyt Application

Recommend Dilutions ELISA 1:5000-10000; Flow Cyt 1:100

Purification Purified from cell culture supernatant by affinity chromatography

Formulation & Reconstitution Lyophilized from sterile PBS, pH 7.4. Normally 5 % – 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.

Scre at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.

This gene encodes a cell surface receptor expresses soon natural filler (NIX) cells, and some T cells) that mediate non-major histocompatibility complex (MHC) restricted killing. The interaction between NK-cell and target cells with this receptor is thought to modulate NK-cell cytolytic activity. Alternatively spliced transcriptor variants encoding different isoforms have been found for this green. Storage & Shipping

Usage Research use only

Necessity used only

Unconjugated

All DIMA recombinant antibodies are genuinely generated by DIMA Biotech. They are all under patent application. Any protein sequencing or engineering attempt is prohibited. We are actively scrutinizing all patent application to ensure no IP infringement. DIMA Disclaimer




Figure 1. ELISA plate pre-coated by 2  $\mu$ g/ml (100  $\mu$ l/well) Human 2B4 protein, mFc-His tagged protein PME100010 can bind Rabbit anti-2B4 monoclonal antibody ( clone: DM69) in a linear range of 1-100 ng/ml.

Email: info@dimabio.com Website: www.dimabio.com

Address: Wuhan institute of Biotechnology B7, Biolake No.666 Gaoxin Road, Wuhan, Hubei, China Telephone: +1 2409940618(USA) /+86-18062749453(China) /+86-400-006-0995(China)






Figure 2. HEK293 cell line transfected with irrelevant protein (A) and human 2B4 (B) were surface stained with Rabbit anti-2B4 monoclonal antibody  $1\mu g/ml$  ( clone: DM69) followed by Alexa 488-conjugated anti-rabbit IgG secondary antibody.




Figure 3. Flow cytometry data of serially titrated Rabbit anti-2B4 monoclonal antibody ( clone: DM69) on THP-1 cells. The Y-axis represents the mean fluorescence intensity (MFI) while the X-axis represents the concentration of IgG used.

Email: info@dimabio.com Website: www.dimabio.com








Figure 4. Affinity ranking of different Rabbit anti-2B4 mAb clones by titration of different concentration onto THP-1 cells. The Y-axis represents the mean fluorescence intensity (MFI) while the X-axis represents the concentration of IgG used.




Figure 5. Flow cytometry analysis of antigen binding of rabbit anti-human 2B4 mAb(DME100069). (A) DME100069 does not bind to CHO-S cells that do not express 2B4. (B) A clear peak shift of DME100069 was seen compared to the control when incubated with 2B4-expressing THP-1 cells, indicating strong binding of DME100069 to 2B4. Antibodies were incubated at 5  $\mu$ g/mL.

ress: Wuhan institute of Email: info@dimabio.com echnology B7, Biolake No.666 Website: www.dimabio.com

