

PRODUCT INFORMATION

Warning . Undefined variable \$hasAttributeValueDescription in Calwawroottmirror.dimablo.com/wp-content\plugins/woocommerce-print-products-public.php on line 2806 bM138 Clone ID

Host Species Rabbit

Anti-5T4 antibody(DM138); Rabbit mAb Description

Delivery In Stock Uniprot ID Q13641 IgG type Rabbit IgG Clonality Monoclonal Reactivity ELISA; Flow Cyt Application

Recommend Dilutions ELISA 1:5000-10000; Flow Cyt 1:100

Purification Purified from cell culture supernatant by affinity chromatography

Formulation & Reconstitution Lyophilized from sterile PBS, pH 7.4. Normally 5 % – 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.

Storage & Shipping

Special instructions reconstitution. Store at 20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature. This gene encodes a leucline-rich transmembrane glycoprotein that may be involved in cell adhesion. The encoded protein is an oncofetal antigen that is specific to trophoblast cells, in adults this protein is highly expressed in many tumor cells and is associated with poor clinical outcome in numerous cancers. Attended splicing in the 5° UTR results in multiple transcript variants that encode the same protein.

Usage Research use only

Unconjugated
All DIMA recombinant antibodies are genuinely generated by DIMA Biotech. They are all under patent application. Any protein engineering attempt is prohibited. We are actively scrutinizing all patent application to ensure no IP infringement. DIMA Disclaimer




Figure 1. ELISA plate pre-coated by 1 μ g/ml (100 μ l/well) Human 5T4 protein, His tagged protein PME100123 can bind Rabbit anti-5T4 monoclonal antibody (clone: DM138) in a linear range of 0.1-16 ng/ml.

Address: Wuhan institute of Biotechnology B7, Biolake No.666 Gaoxin Road, Wuhan, Hubei, China Telephone: +1 2409940618(USA) /+86-18062749453(China) /+86-400-006-0995(China)

Email: info@dimabio.com Website: www.dimabio.com








Figure 2. 5T4 protein is highly expressed on the surface of HEK293 cell membrane. Flow cytometry analysis with Anti-5T4 (DM138) on HEK293 cells transfected with human 5T4 (Blue histogram) or HEK293 transfected with irrelevant protein (Red histogram), and Isotype antibody on HEK293 transfected with irrelevant protein (Orange histogram)





Figure 3. Flow cytometry analysis of antigen binding of anti-human 5T4 mAb(DME100138). (A)DME100138 does weakly bind to Jurkat cells that do not express 5T4. (B) A clear peak shift of DME100138 was seen compared to the control when incubated with 5T4-expressing MCF-7 cells, indicating strong binding of DME100138 to 5T4. Antibodies were incubated at 10 μ g/mL.



Email: info@dimabio.com Website: www.dimabio.com

