

## **PRODUCT INFORMATION**

Warning: Undefined variable ShasAttributeValueDescription in C:\u00e4wroot\u00fcmirror.dimablo.com\u00e4wp-content\plugins\u00e4woocommerce-print-products-public.php on line 2806
274 Clone ID

AGR2 AG-2;hAG-2;HPC8 Host Species

Rabbit Anti-AGR2 antibody(2F4), Rabbit mAb Description Delivery In Stock

Uniprot ID 095994 lgG type Rabbit mAb Clonality Monoclonal Reactivity Human ELISA Applicati

Recommend Dilutions ELISA 1/5000-10000

Purification Purified from cell culture supernatant by affinity chromatography

Formulation & Reconstitution Lyophilized from sterile PBS, pH 7.4. Normally 5 % – 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.

Storage & Shipping

Score at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature. This gene necodes a member of the dissilidies lowerase (PD) family of endoplasmic reticulum (ER) proteins that catalyze protein folding and thiolitisis proteins that catalyze protein folding and thiolitisis of the proteins of the prot Background

Conjugate Unconjugated

## Anti-AGR2 (2F4) mAb ELISA 0.1µg of Human AGR2, hFc tagged protein per well




Figure 1. ELISA plate pre-coated by 1  $\mu$ g/ml (100  $\mu$ l/well) Human AGR2 Protein can bind Rabbit anti-AGR2 monoclonal antibody(clone: 2F4) in a linear range of 1-50 ng/ml.

Address: Wuhan institute of Biotechnology B7, Biolake No.666 Gaoxin Road, Wuhan, Hubei, China Telephone: +1 2409940618(USA) /+86-18062749453(China) /+86-400-006-0995(China)

Email: info@dimabio.com Website: www.dimabio.com

