

## PRODUCT INFORMATION

Warning: Undefined variable ShasAttributeValueDescription in C:\unwwroot\mirror.dimabio.com\wp-content\plugins\woocommerce-printproducts\public\class-woocommerce-print-products-public.php on line 2806
301 Clone ID

AAA;ABETA;ABPP;AD1;APPI;CTFgamma;CVAP;PN-II;PN2;preA4 Synonyme

Host Species

Rabbit Anti-APP antibody(3D7); IgG1 Chimeric mAb Description

Delivery In Stock Uniprot ID P05067

Rabbit/Human Fc chimeric IgG1 IgG type

Clonality Monoclonal Reactivity Human Application Flow Cyt

Recommend Dilutions Flow Cyt 1/100

Purification Purified from cell culture supernatant by affinity chromatography

Formulation & Reconstitution Lyophilized from sterile PBS, pH 7.4. Normally 5 % – 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.

Storage & Shipping

specific instructions of reconstitution.

Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at antibilinent temperature.

This gene encodes a cell surface receptor and transmembrane precursor protein that is cleaved by secretases to form a number of peptides, some of This gene encodes a cell surface receptor and transmembrane precursor protein that is cleaved by secretases to form a number of peptides, some of the protein basis of the amyloid plaques found in the brains of patients with Alzheimer diseases. In addition, two of the peptides are antimicrobial peptides, having been shown to have backerioidal and antifungal activities. Mutations in this gene have been implicated in autosomal dominant Alzheimer disease and cerebroarterial amyloidosis (cerebral amyloid angiopathy). Multiple transcript variants encoding several different isoforms have been foun for this gene. [provided by ReFSeq, Aug 2014]

Usage Research use only

Unconjugated
All DIMA recombinant antibodies are genuinely generated by DIMA Biotech. They are all under patent application. Any protein sequencing or reverse engineering attempt is prohibited. We are actively scrutinizing all patent application to ensure no IP infringement. DIMA Disclaimer




Figure 1. Flow cytometry analysis with Anti-APP (3D7) mAb on HEK293 cells transfected with human APP (Blue histogram) or HEK293 transfected with irrelevant protein (Red histogram).

Address: Wuhan institute of Biotechnology B7, Biolake No.666 Gaoxin Road, Wuhan, Hubei, China Telephone: +1 2409940618(USA) /+86-18062749453(China) /+86-400-006-0995(China)

Email: info@dimabio.com Website: www.dimabio.com



Cat. No. DMC101002







Figure 2. Flow cytometry analysis of antigen binding of anti-human APP mAb(DMC101002). (A) DMC101002 does not bind to CHO-S cells that do not express APP. (B) A clear peak shift of DMC101002 was seen compared to the control when incubated with APP-expressing Siha cells, indicating strong binding of DMC101002 to APP. Antibodies were incubated at 5  $\mu$ g/mL.



Email: info@dimabio.com Website: www.dimabio.com

