

**PRODUCT INFORMATION**

<b>Clone ID</b>	1F2
<b>Target</b>	AXL
<b>Synonyms</b>	AXL;UFO
<b>Host Species</b>	Rabbit
<b>Description</b>	Anti-AXL antibody(1F2), IgG1 Chimeric mAb
<b>Delivery</b>	In Stock
<b>Uniprot ID</b>	P30530
<b>IgG type</b>	Rabbit/Human Fc chimeric IgG1
<b>Clonality</b>	Monoclonal
<b>Reactivity</b>	Human
<b>Applications</b>	WB
<b>Recommended Dilutions</b>	WB 1/1000
<b>Purification</b>	Purified from cell culture supernatant by affinity chromatography
<b>Formulation &amp; Reconstitution</b>	Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.
<b>Storage &amp; Shipping</b>	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
<b>Background</b>	The protein encoded by this gene is a member of the Tyro3-Axl-Mer (TAM) receptor tyrosine kinase subfamily. The encoded protein possesses an extracellular domain which is composed of two immunoglobulin-like motifs at the N-terminal, followed by two fibronectin type-III motifs. It transduces signals from the extracellular matrix into the cytoplasm by binding to the vitamin K-dependent protein growth arrest-specific 6 (Gas6). This gene may be involved in several cellular functions including growth, migration, aggregation and anti-inflammation in multiple cell types. Alternative splicing results in multiple transcript variants of this gene.
<b>Usage</b>	Research use only
<b>Conjugate</b>	Unconjugated
<b>DIMA Disclaimer</b>	All DIMA recombinant antibodies are genuinely generated by DIMA Biotech. They are all under patent application. Any protein sequencing or reverse engineering attempt is prohibited. We are actively scrutinizing all patent application to ensure no IP infringement.






Figure 1. Anti-AXL antibody (SKU# DMC100868) at 1/1000 dilution

Lane : HeLa, whole cell lysate

Secondary : Goat Anti-Rabbit IgG H&L (HRP) at 1/5000 dilution

Predicted band size: 98 kDa  
Observed band size: 128 kDa




Figure 2. Flow cytometry analysis of antigen binding of anti-human AXL mAb(DMC100868).

(A) DMC100868 does not bind to CHO-S cells that do not express AXL.

(B) A clear peak shift of DMC100868 was seen compared to the control when incubated with AXL-expressing HeLa cells, indicating strong binding of DMC100868 to AXL. Antibodies were incubated at 5  $\mu$ g/mL.

