

PRODUCT INFORMATION

Warning: Undefined variable ShasAttributeValueDescription in C:\u00e4wroot\u00e4mirror.dimablo.com\u00e4wp-content\plugins\u00e4woocommerce-print-products\u00e4public\u00e4public\u00e4pub on line 2806 125 Clone ID

AXL;UFO Synonyme Host Species Rabbit

Anti-AXL antibody(1F2), IgG1 Chimeric mAb Description

Delivery In Stock Uniprot ID P30530

Rabbit/Human Fc chimeric IgG1 lgG type

Clonality Monoclonal Reactivity Human Application WB Recommend Dilutions WB 1/1000

Purification Purified from cell culture supernatant by affinity chromatography

Formulation & Reconstitution Lyophilized from sterile PBS, pH 7.4. Normally 5 % – 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.

Storage & Shipping

specific instructions of reconstitution.

Store at -20°C to .80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.

The protein encoded by this gene is a member of the Tyro-1A-vike (TAB) receptor tyrosine kinase subfamily. The encoded protein possesses an extracellular domain which is composed of two immunoglobulin-like motifs at the N-terminal, followed by two fibronectin type-III motifs. It transduces signals from the extracellular matrix into the cytoplasm by binding to the vitamin K-dependent protein growth are specific 6 (Gas5). This gene may be involved in several cellular functions including growth, migration, aggregation and anti-inflammation in multiple cell types. Alternative splicing Research use only Background

Usage

Coniugate

All DIMA recombinant antibodies are genuinely generated by DIMA Biotech. They are all under patent application. Any protein sequencing or reverse engineering attempt is prohibited. We are actively scrutinizing all patent application to ensure no IP infringement. DIMA Disclaime

250 kDa -

130 kDa -

100 kDa -

70 kDa -

55 kDa -

35 kDa -

25 kDa -

Figure 1.Anti-AXL antibody (SKU# DMC100868) at 1/1000 dilution

Lane: Hela, whole cell lysate

Secondary: Goat Anti-Rabbit IgG H&L (HRP) at 1/5000 dilution

Predicted band size: 98 kDa Observed band size: 128 kDa

/+86-400-006-0995(China)

Address: Wuhan institute of Biotechnology B7, Biolake No.666 Gaoxin Road, Wuhan, Hubei, China Telephone: +1 2409940618(USA) /+86-18062749453(China)

Email: info@dimabio.com Website: www.dimabio.com








Figure 2. Flow cytometry analysis of antigen binding of anti-human AXL mAb(DMC100868). (A) DMC100868 does not bind to CHO-S cells that do not express AXL. (B) A clear peak shift of DMC100868 was seen compared to the control when incubated with AXL-expressing Hela cells, indicating strong binding of DMC100868 to AXL. Antibodies were incubated at 5 μ g/mL.

Address: Wuhan institute of Biotechnology B7, Biolake No.666 Gaoxin Road, Wuhan, Hubei, China Telephone: +1 2409940618(USA) /+86-18062749453(China) /+86-400-006-0995(China)

Email: info@dimabio.com Website: www.dimabio.com

