

PRODUCT INFORMATION

Warning: Undefined variable ShasAttributeValueDescription in C:\u00e4wroot\u00e4mirror.dimablo.com\u00e4wp-content\plugins\u00e4woocommerce-print-products\u00e4public\u00e7class-woocommerce-print-products-public.php on line 2806
bM66 Clone ID

CD86; B7-2; B70; CD28LG2; LAB72; MGC34413

Host Species Rabbit

Anti-B7-2 antibody(DM86); Rabbit mAb Description 3~4 weeks

Delivery P42081 Uniprot ID IgG type Rabbit IgG Clonality Monoclonal Reactivity Application ELISA; Flow Cyt

Recommend Dilutions ELISA 1:5000-10000; Flow Cyt 1:100

Purification Purified from cell culture supernatant by affinity chromatography

Formulation & Reconstitution Lyophilized from sterile PBS, pH 7.4. Normally 5 % – 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.

Storage & Shipping

specific instructions of reconstitution.

Store at 2-0°C to .80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.

This gene encodes a type I membrane protein that is a member of the immunoglobulin superfamily. This protein is expressed cells; and it is the ligand for two proteins at the cell surface of T cells; CD28 antigen and cytotoxic T-lymphocyte-associated protein with CD28 antigen is a costimulatory signal for activation of the T-cell. Binding of this protein with cytoric T-lymphonegatively regulates T-cell activation and diminishes the immune response. Alternative splicing results in several transcript v isoforms.

Research use only

Usage

Coniugate

Background

All DIMA recombinant antibodies are genuinely generated by DIMA Biotech. They are all under patent application. Any protein sequencing or reverse engineering attempt is prohibited. We are actively scrutinizing all patent application to ensure no IP infringement. DIMA Disclaime




Figure 1. ELISA plate pre-coated by 2 μ g/ml (100 μ l/well) Human B7-2 protein, mFc-His tagged protein PME100034 can bind Rabbit anti-B7-2 monoclonal antibody (clone: DM86) in a linear range of 1-100 ng/ml.

Address: Wuhan institute of Biotechnology B7, Biolake No.666 Gaoxin Road, Wuhan, Hubei, China Telephone: +1 2409940618(USA) /+86-18062749453(China) /+86-400-006-0995(China)

Email: info@dimabio.com Website: www.dimabio.com








Figure 2. HEK293 cell line transfected with irrelevant protein (A) and human B7-2 (B) were surface stained with Rabbit anti-B7-2 monoclonal antibody 1µg/ml (clone: DM86) followed by Alexa 488-conjugated anti-rabbit IgG secondary antibody.





Figure 3. Flow cytometry analysis of antigen binding of rabbit anti-human B7-2 mAb(DME100086). (A) DME100086 does not bind to 293T cells that do not express B7-2. (B) A clear peak shift of DME100086 was seen compared to the control when incubated with B7-2-expressing Daudi cells, indicating strong binding of DME100086 to B7-2. Antibodies were incubated at 5 $\mu g/mL$







Cat. No. DME100086






Figure 4. Flow cytometry analysis of antigen binding of rabbit anti-human B7-2 mAb(DME100086). (A) DME100086 does not bind to 293T cells that do not express B7-2. (B) A clear peak shift of DME100086 was seen compared to the control when incubated with B7-2-expressing Raji cells, indicating strong binding of DME100086 to B7-2. Antibodies were incubated at 5 μ g/mL.

