## **PRODUCT INFORMATION**

Warning: Undefined variable ShasAttributeValueDescription in C:\www.root\mirror.dimablo.com\wp-content\plugins\woocommerce-print-products\public\class-woocommerce-print-products-public.php on line 2806 AMG557, AMG-557, MEDI-5872

Conjugate

Unconjugated ICOSLG;B7-H2;B7RP-1;B7RP1;CD275;GL50;ICOS-L;ICOSL;LICOS;ICOS ligand Synonyms

Applications Flow Cyt Recommend Dilutions Flow Cyt 1:100

Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution. Formulation & Reconstitution

Human IgG2 - Kappa IgG type Reactivity Target B7-H2 Uniprot ID


Anti-B7-H2(prezalumab biosimilar) mAb Description

Delivery

In Stock
Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.

Background Research grade biosimilar. Not for use in therapeutic or diagnostic procedures for humans or animals

Usage



**Figure 1.** Flow cytometry analysis with 15  $\mu$ g/mL Anti-B7-H2 (prezalumab biosimilar) mAb (BME100081) on HEK293 cells transfected with Human B7H2 protein (Blue histogram) or HEK293 transfected with irrelevant protein (Red histogram).

Address: Wuhan institute of Biotechnology B7, Biolake No.666 Gaoxin Road, Wuhan, Hubei, China Telephone: +1 2409940618(USA) /+86-18062749453(China) /+86-400-006-0995(China)

Email: info@dimabio.com Website: www.dimabio.com

