

PRODUCT INFORMATION

Warning: Undefined variable ShasAttributeValueDescription in C:\www.root\mirror.dimablo.com\wp-content\plugins\woocommerce-print-products\public\class-woocommerce-print-products-public.php on line 2806 bMC299 Clone ID

B7-H5; B7H5; C10orf54; DD1alpha; Dies1; Gl24; PD-1H; PP2135; SISP1; VISTA Synonyme

Host Species

Anti-B7-H5 antibody(DMC299); IgG1 Chimeric mAb Description

Delivery In Stock Uniprot ID

Q9H7M9 Rabbit/Human Fc chimeric IgG1 lgG type

Clonality Monoclonal Reactivity Human Application Flow Cyt Recommend Dilutions Flow Cyt 1:100

Purification Purified from cell culture supernatant by affinity chromatography

Formulation & Reconstitution Lyophilized from sterile PBS, pH 7.4. Normally 5 % – 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.

Store at 2-0°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature. In the store of the s Storage & Shipping Background

Conjugate Unconjugated

All DIMA recombinant antibodies are genuinely generated by DIMA Biotech. They are all under patent application. Any protein sequencing or reverse engineering attempt is prohibited. We are actively scrutinizing all patent application to ensure no IP infringement. DIMA Disclaimer




Figure 1. Flow cytometry analysis with Anti-B7H5 (DMC299) on HEK293 cells transfected with human B7H5 (Blue histogram) or HEK293 transfected with irrelevant protein (Red histogram).

Address: Wuhan institute of Biotechnology B7, Biolake No.666 Gaoxin Road, Wuhan, Hubei, China Telephone: +1 2409940618(USA) /+86-18062749453(China) /+86-400-006-0995(China)

Email: info@dimabio.com Website: www.dimabio.com



Cat. No. DMC100299






Figure 2. Flow cytometry analysis of antigen binding of anti-human B7-H5 mAb(DMC100299). (A) DMC100299 does not bind to 293T cells that do not express B7-H5. (B) A clear peak shift of DMC100299 was seen compared to the control when incubated with B7-H5-expressing THP-1 cells, indicating strong binding of DMC100299 to B7-H5. Antibodies were incubated at 5 μ g/mL.





