

## PRODUCT INFORMATION

iable \$hasAttributeValueDescription in C:\wwwroot\mirror.dimabio.com\wp-content\plugins\woocommerce-print-woocommerce-print-products-public.php on line 2806 Clone ID

PVR; FLJ25946; PVS; CD155; TAGE4; HVED; NECL5

Host Species Rabbit

Anti-CD155 antibody(DM146); Rabbit mAb Description Delivery In Stock

Uniprot ID P15151 IgG type Rabbit IgG Clonality Monoclonal Reactivity Human ELISA; Flow Cyt

Application

Background

Recommend Dilutions ELISA 1:5000-10000; Flow Cyt 1:100

Purification Purified from cell culture supernatant by affinity chromatography

Formulation & Reconstitution Lyophilized from sterile PBS, pH 7.4. Normally 5 % – 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.

Scre at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at a mishient temperature in the protein benefit of the primate lineage; and serves as a cellular receptor for poliovirus in the first step of poliovirus replication. Multiple transcript variants encoding different isoforms have been found for this gene. Storage & Shipping

Usage Conjugate Research use only Unconjugated

All DIMA recombinant antibodies are genuinely generated by DIMA Biotech. They are all under patent application. Any protein sequencing or reverse engineering attempt is prohibited. We are actively scrutinizing all patent application to ensure no IP infringement.




Figure 1. ELISA plate pre-coated by 1  $\mu$ g/ml (100  $\mu$ l/well) Human CD155 protein, mFc tagged protein PME100027 can bind Rabbit anti-CD155 monoclonal antibody (clone: DM146) in a linear range of 5-100 ng/ml.

Address: Wuhan institute of Biotechnology B7, Biolake No.666 Gaoxin Road, Wuhan, Hubei, China Telephone: +1 2409940618(USA) /+86-18062749453(China) /+86-400-006-0995(China)

Email: info@dimabio.com Website: www.dimabio.com








Figure 2. CD155 protein is highly expressed on the surface of HEK293 cell membrane. Flow cytometry analysis with Anti-CD155 (DM146) on HEK293 cells transfected with human CD155 (Blue histogram) or HEK293 transfected with irrelevant protein (Red histogram), and Isotype antibody on HEK293 transfected with irrelevant protein (Orange histogram).




Figure 3. Flow cytometry analysis of antigen binding of rabbit anti-human CD155 mAb(DME100146). (A) DME100146 does not bind to CHO-S cells that do not express CD155. (B) A clear peak shift of DME100146 was seen compared to the control when incubated with CD155-expressing Daudi cells, indicating strong binding of DME100146 to CD155. Antibodies were incubated at 5  $\mu$ g/ml.



Email: info@dimabio.com Website: www.dimabio.com

