

PRODUCT INFORMATION

| | |
|---|--|
| Clone ID | 1D4 |
| Target | CD205 |
| Synonyms | LY75;CLEC13B;DEC-205;GP200-MR6;LY-75 |
| Host Species | Rabbit |
| Description | Anti-CD205 antibody(1D4), IgG1 Chimeric mAb |
| Delivery | In Stock |
| Uniprot ID | O60449 |
| IgG type | Rabbit/Human Fc chimeric IgG1 |
| Clonality | Monoclonal |
| Reactivity | Human |
| Applications | Flow Cyt |
| Recommended Dilutions | Flow Cyt 1/100 |
| Purification | Purified from cell culture supernatant by affinity chromatography |
| Formulation & Reconstitution | Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution. Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature. |
| Storage & Shipping | Acts as an endocytic receptor to direct captured antigens from the extracellular space to a specialized antigen-processing compartment (By similarity). Causes reduced proliferation of B-lymphocytes.[UniProtKB/Swiss-Prot Function] |
| Background | |
| Usage | Research use only |
| Conjugate | Unconjugated |
| DIMA Disclaimer | All DIMA recombinant antibodies are genuinely generated by DIMA Biotech. They are all under patent application. Any protein sequencing or reverse engineering attempt is prohibited. We are actively scrutinizing all patent application to ensure no IP infringement. |






Figure 1. Flow cytometry analysis with 1µg/mL Anti-CD205 (1D4) mAb on Expi293 cells transfected with human CD205 (Blue histogram) or Expi293 transfected with irrelevant protein (Red histogram).




Figure 2. Flow cytometry analysis of antigen binding of anti-human CD205 mAb(DMC101114).

(A) DMC101114 does not bind to Jurkat cells that do not express CD205.

(B) A clear peak shift of DMC101114 was seen compared to the control when incubated with CD205-expressing Raji cells, indicating strong binding of DMC101114 to CD205. Antibodies were incubated at 5 µg/mL.

