

PRODUCT INFORMATION

Warning: Undefined variable ShasAttributeValueDescription in C:\www.root\mirror.dimablo.com\wp-content\plugins\woocommerce-print-products\public\class-woocommerce-print-products-public.php on line 2806
bMS8 Clone ID

CD27; TNFRSF7; S152; T14; Tp55

Host Species Rabbit

Anti-CD27 antibody(DM58); Rabbit mAb Description In Stock

Delivery Uniprot ID P26842 lgG type Rabbit IgG Clonality Monoclonal Reactivity

Application

ELISA; Flow Cyt Recommend Dilutions ELISA 1:5000-10000; Flow Cyt 1:100

Purification Purified from cell culture supernatant by affinity chromatography

Formulation & Reconstitution Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.

Storage & Shipping

specific instructions of reconstitution.

Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.

The protein encoded by this gene is a member of the TIM-receptor superfamily. This receptor is required for generation and long-term maintenance of cell immunity. It binds to ligand CD70; and plays a key role in regulating B-cell activation and immunoglobulin synthesis. This receptor transduces signals that lead to the activation of NF-kappaB and MAPKB;IMA. Adaptor proteins TRAP2 and TRAP5 have been shown to mediate the signaling process of this receptor. CD27-binding protein (SIVA); a proapoptotic protein; can bind to this receptor and is thought to play an important role in the apoptosis induced by this receptor.

Research use only Background

Usage Coniugate

All DIMA recombinant antibodies are genuinely generated by DIMA Biotech. They are all under patent application. Any protein sequencing or reverse engineering attempt is prohibited. We are actively scrutinizing all patent application to ensure no IP infringement. DIMA Disclaime




Figure 1. HEK293 cell line transfected with irrelevant protein (A) and human CD27 (B) were surface stained with Rabbit anti-CD27 monoclonal antibody 1µg/ml (clone: DM58) followed by Alexa 488-conjugated anti-rabbit IgG secondary antibody.

Email: info@dimabio.com Website: www.dimabio.com










Figure 2. Flow cytometry data of serially titrated Rabbit anti-CD27 monoclonal antibody (clone: DM58) on Raji cells. The Y-axis represents the mean fluorescence intensity (MFI) while the X-axis represents the concentration of IgG used.




Figure 3. Affinity ranking of different Rabbit anti-CD27 mAb clones by titration of different concentration onto Raji cells. The Y-axis represents the mean fluorescence intensity (MFI) while the X-axis represents the concentration of IgG used.

Email: info@dimabio.com Website: www.dimabio.com



Cat. No. DME100058






Figure 4. Flow cytometry analysis of antigen binding of rabbit anti-human CD27 mAb(DME100058).

(A) DME100058 does not bind to 293T cells that do not express CD27. (B) A clear peak shift of DME100058 was seen compared to the control when incubated with CD27-expressing Raji cells, indicating strong binding of DME100058 to CD27. Antibodies were incubated at 2 μ g/mL.

Address: Wuhan institute of Biotechnology B7, Biolake No.666 Gaoxin Road, Wuhan, Hubei, China Telephone: +1 2409940618(USA) /+86-18062749453(China) /+86-400-006-0995(China)

Email: info@dimabio.com Website: www.dimabio.com

