

PRODUCT INFORMATION

Common Name	Warning: Undefined variable \$hasAttributeValueDescription in C:\wwwroot\mirror.dimabio.com\wp-content\plugins\woocommerce-print-products\publicclass-woocommerce-print-products-public.php on line 2806 IF5, CDX-1127
Synonyms	Warning: Undefined variable \$hasAttributeValueDescription in C:\wwwroot\mirror.dimabio.com\wp-content\plugins\woocommerce-print-products\publicclass-woocommerce-print-products-public.php on line 2806 CD27, TNFRSF7, S152, T14, Tp55
Conjugate	Unconjugated
Applications	ELISA; Flow Cyt
Recommended Dilutions	ELISA 1:5000-10000; Flow Cyt 1:100
Formulation & Reconstitution	Lyophilized from sterile PBS, pH 7.4. Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.
Host Species	Homo sapiens
IgG type	Human IgG1 - kappa
Reactivity	Human
Target	CD27
Uniprot ID	P26842
Description	Anti-CD27 (varlilumab biosimilar) mAb
Delivery	In Stock
Yefei_Storage	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
Background	Research grade biosimilar. Not for use in therapeutic or diagnostic procedures for humans or animals.
Usage	Research use only

Human CD27, hFc Tagged protein ELISA

0.2 µg of CD27, hFc Tagged protein per well




Figure 1. ELISA plate pre-coated by 2 µg/ml (100 µL/well) Human CD27, hFc tagged protein (PME100475) can bind Anti-CD27 Neutralizing antibody in a linear range of 3.2-80.0 ng/ml. In order to specifically detect BME100018, mouse anti-human Fab-specific antibody was used as detection antibody.



Anti-CD27 (varlilumab biosimilar) mAb ELISA

0.2 μ g of CD27, mFc-His Tagged protein per well




Figure 2. ELISA plate pre-coated by 2 μ g/ml (100 μ l/well) Human CD27, mFc-His tagged protein ([getskuurl sku="PME100013"]) can bind Anti-CD27 Neutralizing antibody in a linear range of 0.64-16.0 ng/ml.




Figure 3. HEK293 cell line transfected with irrelevant protein (**B**) and human CD27 protein (**A**) were surface stained with anti-CD27 neutralizing antibody 1 μ g/ml (varlilumab) followed by Alexa 488-conjugated anti-human IgG secondary antibody.

