

PRODUCT INFORMATION

Common Name	Warning: Undefined variable \$hasAttributeValueDescription in C:\wwwroot\mirror\dimabio.com\wp-content\plugins\woocommerce-print-products\public\class-woocommerce-print-products-public.php on line 2806 MK-5890, MK5890
Conjugate	Unconjugated
Synonyms	CD27, TNFRSF7, S152, T14, Tp55
Applications	ELISA, Flow Cyt
Recommended Dilutions	ELISA 1:5000-10000, Flow Cyt 1:100
Formulation & Reconstitution	Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.
Host Species	Chimeric Humanized
IgG type	Human IgG1(E356D,M358L) - kappa
Reactivity	Human
Target	CD27
Uniprot ID	P26842
Description	Anti-CD27(boserolimab biosimilar) mAb
Delivery	In Stock
Yefei Storage	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized antibodies are shipped at ambient temperature.
Background	Research grade biosimilar. Not for use in therapeutic or diagnostic procedures for humans or animals.
Usage	Research use only

Anti-CD27(boserolimab biosimilar) mAb ELISA

0.2 µg of Human CD27, mFc-His tagged protein per well




Figure 1. ELISA plate pre-coated by 2 µg/mL (100 µL/well) Human CD27 Protein, mFc-His Tag (PME100013) can bind Anti-CD27(boserolimab biosimilar) mAb (BME100439) in a linear range of 1.6–8 ng/mL.






Figure 2. Flow cytometry analysis with 1 μ g/mL Anti-CD27(boserolimab biosimilar) mAb (BME100439) on HEK293 cells transfected with Human CD27 protein (Blue histogram) or HEK293 transfected with irrelevant protein (Red histogram).

DIMABIO CONFIDENTIAL

