

## **PRODUCT INFORMATION**

Warning: Undefined variable \$hasAttributeValueDescription in C:\www.root\mirror.dimablo.com\wp-content\plugins\woocommerce-print-products\public.php on line 2806 MK-5890, MK-5890.

Conjugate

Unconjugated CD27, TNFRSF7, S152, T14, Tp55 Synonyms Applications ELISA, Flow Cyt

Recommend Dilutions ELISA 1:5000-10000, Flow Cyt 1:100

Lyophilized from sterile PBS, pH 7.4. Normally 5 % – 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution. Formulation & Reconstitution

Human IgG1(E356D,M358L) - kappa IgG type

Reactivity Target CD27 Uniprot ID

Anti-CD27(boserolimab biosimilar) mAb Description

Delivery


In Stock
Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized antibodies are shipped at ambient temperature.

Background Research grade biosimilar. Not for use in therapeutic or diagnostic procedures for humans or animals

Usage

## Anti-CD27(boserolimab biosimilar) mAb ELISA

0.2 μg of Human CD27, mFc-His tagged protein per well



Anti-CD27(boserolimab biosimilar) mAb (ng/mL)

Figure 1. ELISA plate pre-coated by 2  $\mu$ g/mL (100  $\mu$ L/well) Human CD27 Protein, mFc-His Tag (PME100013) can bind Anti-CD27(boserolimab biosimilar) mAb (BME100439) in a linear range of 1.6–8 ng/mL.



Email: info@dimabio.com Website: www.dimabio.com








Figure 2. Flow cytometry analysis with 1  $\mu$ g/mL Anti-CD27(boserolimab biosimilar) mAb (BME100439) on HEK293 cells transfected with Human CD27 protein (Blue histogram) or HEK293 transfected with irrelevant protein (Red histogram).

Email: info@dimabio.com Website: www.dimabio.com

