

PRODUCT INFORMATION

Warning: Undefined variable ShasAttributeValueDescription in C:\www.root\mirror.dimablo.com\wp-content\plugins\woocommerce-print-products\public.class-woocommerce-print-products-public.php on line 2806 bM193 Clone ID

B220; CD45; CD45R; GP180; L-CA; LCA; LY5; T200 Synonyme

Host Species Rabbit

Anti-CD45 antibody(DM193); Rabbit mAb Description In Stock

Delivery Uniprot ID P08575 IgG type Rabbit IgG Clonality Monoclonal Reactivity Human Application ELISA; Flow Cyt

Recommende Dilutions ELISA 1:5000-10000; Flow Cyt 1:100

Purification Purified from cell culture supernatant by affinity chromatography

Formulation & Reconstitution Lyophilized from sterile PBS, pH 7.4. Normally 5 % – 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.

Storage & Shipping

specific instructions of reconstitution.

Store at 20°C to 80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.

The protein encoded by this gene is a member of the protein in vorsine phosphatase (PTP) family. PTPs are known to be signaling molecules that regulate a variety of cellular processes including cell growth; differentiation; milosis; and oncogenic transformation. This PTP contains an extracellular domain; a single transmembrane segment and two tandem intracytoplasmic catalytic domains; and thus is classified as expector type PTP. This PTP has been shown to be an essential regulator of T- and B-cell antigen receptor signaling. It functions through either direct interaction with components of the antigen receptor or by activating various STc family kinases required for the antigen receptor signaling. This PTP also suppresses JAK kinases, and thus functions as a regulator of cytokine receptor signaling. Alternatively spliced transcripts variants of this gene; which encode distinct sloftens; have been reported.

Conjugate Unconjugated

Background

DIMA Disclaimer All DIMA recombinant antibodies are genuinely generated by DIMA Biotech. They are all under patent application. Any protein sequencing or reverse engineering attempt is prohibited. We are actively scrutinizing all patent application to ensure no IP infringement.




Figure 1. Flow cytometry analysis with Anti-CD45 (DM193) on HEK293 cells transfected with human CD45 (Blue histogram) or HEK293 transfected with irrelevant protein (Red histogram).

Address: Wuhan institute of Biotechnology B7, Biolake No.666 Gaoxin Road, Wuhan, Hubei, China Telephone: +1 2409940618(USA) /+86-18062749453(China)

/+86-400-006-0995(China)

Email: info@dimabio.com Website: www.dimabio.com








Figure 2. Flow cytometry analysis of antigen binding of rabbit anti-human CD45 mAb(DME100193). (A) DME100193 does not bind to CHO-S cells that do not express CD45. (B) A clear peak shift of DME100193 was seen compared to the control when incubated with CD45-expressing Jurkat cells, indicating strong binding of DME100193 to CD45. Antibodies were incubated at 5 μ g/mL.




Figure 3. Flow cytometry analysis of antigen binding of rabbit anti-human CD45 mAb(DME100193). (A) DME100193 does not bind to CHO-S cells that do not express CD45. (B) A clear peak shift of DME100193 was seen compared to the control when incubated with CD45-expressing THP-1 cells, indicating strong binding of DME100193 to CD45. Antibodies were incubated at 5 μ g/mL.

Email: info@dimabio.com Website: www.dimabio.com

