

## **PRODUCT INFORMATION**

**Common Name** Hu5F9-G4

**Synonyms** CD47;MER6;IAP **Applications** ELISA; Flow Cyt

Recommended

**Dilutions** 

ELISA 1:5000-10000; Flow Cyt 1:100

Lyophilized from sterile PBS, pH 7.4. Normally 5 % – 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis Formulation & Reconstitution

for specific instructions of reconstitution.

**Host Species** Chimeric Humanized

IgG type IgG4 Reactivity Human **CD47 Target Uniprot ID** Q08722

**Description** Anti-CD47 (magrolimab biosimilar, IgG4) mAb

**Delivery** In Stock

> Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not

intended for use within a month, aliquot and store Storage & Shipping

at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient

temperature.

Research grade biosimilar. Not for use in

therapeutic or diagnostic procedures for humans **Background** 

or animals.

Usage Research use only

Conjugate Unconjugated

> All DIMA recombinant antibodies are genuinely generated by DIMA Biotech. They are all under patent application. Any protein sequencing or reverse engineering attempt is prohibited. We are

> > Email: info@dimabio.com Website: www.dimabio.com

**DIMA Disclaimer** 

actively scrutinizing all patent application to ensure no IP infringement.





## Human CD47, mFc-His Tagged protein ELISA

0.2 µg of Human CD47, mFc-His tagged protein per well




Figure 1. ELISA plate pre-coated by 2  $\mu$ g/ml (100  $\mu$ l/well) Human CD47, mFc-His tagged protein PME100008 can bind Anti-CD47 (Neutralizing antibody clone Hu5F9-G4) in a linear range of 0.061-1.606 ng/ml.




Figure 2. HEK293 cell line transfected with irrelevant protein (B) and human CD47(A) were surface stained with anti-CD47 neutralizing antibody 1µg/ml (magrolimab) followed by Alexa 488-conjugated anti-human IgG secondary antibody.



