

PRODUCT INFORMATION

Common Name BB-10901, IMGN901, huN901-DM1

Synonyms MSK39;NCAM **Applications** ELISA; Flow Cyt

Recommended **Dilutions**

ELISA 1:5000-10000; Flow Cyt 1:100

Formulation & Reconstitution Lyophilized from sterile PBS, pH 7.4. Normally 5 % – 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.

Host Species Humanized

IgG type lgG1 Reactivity Human **CD56 Target Uniprot ID** P13591

Description Anti-CD56(lorvotuzumab biosimilar) mAb

Delivery In Stock

> Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not

intended for use within a month, aliquot and store Storage & Shipping

at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient

temperature.

Research grade biosimilar. Not for use in

therapeutic or diagnostic procedures for humans **Background**

or animals.

Usage Research use only

Conjugate Unconjugated

> All DIMA recombinant antibodies are genuinely generated by DIMA Biotech. They are all under patent application. Any protein sequencing or reverse engineering attempt is prohibited. We are

> > Email: info@dimabio.com Website: www.dimabio.com

DIMA Disclaimer

actively scrutinizing all patent application to ensure no IP infringement.





Anti-CD56 (Iorvotuzumab biosimilar) mAb ELISA

0.2 μg of Human CD56, His tagged protein per well




Figure 1. ELISA plate pre-coated by 2 μ g/mL (100 μ L/well) Human CD56 Protein, His Tag ([getskuurl sku="PME100194"]) can bind Anti-CD56 Neutralizing antibody (BME100102) in a linear range of 0.51-41.15 ng/mL.

Anti-CD56 (lorvotuzumab biosimilar) mAb ELISA

0.2 μg of Human CD56, hFc tagged protein per well




Figure 2. ELISA plate pre-coated by 2 μ g/mL (100 μ L/well) Human CD56 Protein, hFc Tag (PME100540) can bind Anti-CD56 Neutralizing antibody (BME100102) in a linear range of 1.52–123.46 ng/mL. In order to specifically detect BME100102, mouse anti-human Fab-specific antibody was used as detection antibody.

Address: Wuhan institute of Biotechnology B7, Biolake No.666 Gaoxin Road, Wuhan, Hubei, China Telephone: +1 2409940618(USA) /+86-18062749453(China) /+86-400-006-0995(China) Email: info@dimabio.com Website: www.dimabio.com






Figure 3. Flow cytometry analysis with 1 μ g/mL Anti-CD56 (lorvotuzumab biosimilar) mAb (BME100102) on Expi293 cells transfected with Human CD56 protein (Blue histogram) or Expi293 transfected with irrelevant protein (Red histogram).



Email: info@dimabio.com Website: www.dimabio.com

