

PRODUCT INFORMATION

Warning: Undefined variable ShasAttributeValueDescription in C:\www.root\mirror.dimablo.com\wp-content\plugins\woocommerce-print-products\public.class-woocommerce-print-products-public.php on line 2806 bMC441 Clone ID

AIM; API6; CT-2; hAIM; PRO229; SP-ALPHA; Spalpha Synonyme

Host Species Rabbit

Anti-CD5L antibody(DMC441); IgG1 Chimeric mAb Description

Delivery In Stock Uniprot ID

043866 Rabbit/Human Fc chimeric IgG1 lgG type

Clonality Monoclonal Reactivity Human Flow Cyt Application Recommende Dilutions Flow Cyt 1:100

Purification Purified from cell culture supernatant by affinity chromatography

Formulation & Reconstitution Lyophilized from sterile PBS, pH 7.4. Normally 5 % – 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.

Storage & Shipping

Lyophilized from Stenie PS, pn 7.4. Normany 3 % - 5% deninuse is adviced as protections betwee syspamics in rease see certained.

Store at 20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing), Lyophilized proteins are shipped at ambient temperature.

repeated freezing and thawing), Lyophilized proteins are shipped at ambient temperature.

rechanisms in inflammatory responses; such as infection or atherosclerosis, Able to inhibit lipid droplet size in adjocytes. Following incorporation into mature adipocytes via CD36-mediated endocytosis; associates with cytosolic RASN, inhibiting flaty acid synthase such sets of the degradation of triacy(plycerols into glycerol and free fatty acids (FFA). CD51-induced lipolysis occurs with progression of obesity: participates in obesity associated inflammatory macrophages into adjocs tissues, a cause of insistive via deliant of inflammatory macrophages into adjocs tissues, a cause of insisting observing the such as a key requisitor of metabolic switch in T-helper Th17 cells, Requisites the expression of pro-inflammatory genes in Th17. Cells in the control of the

Research use only Unconjugated

Conjugate

DIMA Disclaime All DIMA recombinant antibodies are genuinely generated by DIMA Biotech. They are all under patent application. Any protein sequencing or reverse engineering attempt is prohibited. We are actively scrutinizing all patent application to ensure no IP infringement.




Figure 1. Flow cytometry analysis with Anti-CD5L (DMC441) on HEK293 cells transfected with human CD5L (Blue histogram) or HEK293 transfected with irrelevant protein (Red histogram).

Address: Wuhan institute of Biotechnology B7, Biolake No.666 Gaoxin Road, Wuhan, Hubei, China Telephone: +1 2409940618(USA) /+86-18062749453(China) /+86-400-006-0995(China)

Email: info@dimabio.com Website: www.dimabio.com

