

## **PRODUCT INFORMATION**

Warning: Undefined variable ShasAttributeValueDescription in C:\www.root\mirror.dimablo.com\wp-content\plugins\woocommerce-print-products\public\class-woocommerce-print-products-public.php on line 2806 bMc425 Clone ID

DMC-12-2 CD63 antigen; Granulophysin; LAMP-3; Limp1; Melanoma-associated antigen ME491; OMA81H; Ocular melanoma-associated antigen; Tetraspanin-30; Tspan-30 Synonyme

Host Species Rabbit

Anti-CD63 antibody(DMC425); IgG1 Chimeric mAb In Stock

Delivery

IgG type

Uniprot ID P08962 Rabbit/Human Fc chimeric IgG1

Monoclonal Reactivity Human Annlicati

Recommende Dilutions

Formulation & Reconstitution

Lyophilized from sterile PBS, pH 7.4. Normally 5 % – 9% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.

Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoi repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.

The protein encoded by this gene is a member of the transmembrane 4 superfamily; also known as the tetraspanin family. Most of these members a cell-surface proteins that are characterized by the presence of four hydrophobic domains. The proteins mediate signal transduction events that play role in the regulation of cell development; activation; growth and modifyl. The encoded protein is a cell surface deprotein that is known to comple with integrins. It may function as a blood platelet activation marker. Dericiency of this protein is associated with Hermanisty-Puddak syndrome. Also til gene has been associated with unmor progression. Attenative Splicing results in multiple transcript variants encoding different protein is forms.

Usage Research use only

Conjugate All DIMA recombinant antibodies are genuinely generated by DIMA Biotech. They are all under patent application. Any protein sequencing or reverse engineering attempt is prohibited. We are actively scrutinizing all patent application to ensure no IP infringement. DIMA Disclaim




Figure 1. CD63 protein is highly expressed on the surface of HEK293 cell membrane. Flow cytometry analysis with Anti-CD63 (DMC425) on HEK293 cells transfected with human CD63 (Blue histogram) or HEK293 transfected with irrelevant protein (Red histogram), and Isotype antibody on HEK293 transfected with irrelevant protein (Orange histogram).

Address: Wuhan institute of Biotechnology B7, Biolake No.666 Gaoxin Road, Wuhan, Hubei, China Telephone: +1 2409940618(USA) /+86-18062749453(China) /+86-400-006-0995(China)

Email: info@dimabio.com Website: www.dimabio.com



Cat. No. DMC100425






Figure 2. Flow cytometry analysis of antigen binding of anti-human CD63 mAb(DMC100425). (A) DMC100425 does not bind to CHO-S cells that do not express CD63. (B) A clear peak shift of DMC100425 was seen compared to the control when incubated with CD63-expressing THP-1 cells, indicating strong binding of DMC100425 to CD63. Antibodies were incubated at 5  $\mu$ g/mL.



