

## **PRODUCT INFORMATION**

\$hasAttributeValueDescription in C:\wwwroot\mirror.dimabio.com\wp-content\plugins\woocommerce-printcommerce-print-products-public.php on line 2806

Conjugate

Unconjugated NT5E, 5'-Nucleotidase, 5'-NT, NT5, NTE Synonyms

Applications Flow Cyt Recommend Dilutions Flow Cyt 1:100

Lyophilized from sterile PBS, pH 7.4. Normally 5 % – 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution. Formulation & Reconstitution

Human IgG2(C223S,Q274K,F296Y,F300Y,V309L,G327A,A330S,P331S,T339A,S378A,M397V) - Kappa IgG type

Reactivity Target CD73 Uniprot ID

Anti-CD73(BMS 986179) mAb Description

Delivery

In Stock
Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized antibodies are shipped at ambient temperature.

Background Research grade biosimilar. Not for use in therapeutic or diagnostic procedures for humans or animals

Usage




Figure 1. Flow cytometry analysis with  $15\mu g/mL$  Anti-CD73(BMS 986179) mAb (BME100218) on HEK293 cells transfected with Human CD73 protein (Blue histogram) or HEK293 transfected with irrelevant protein (Red histogram).

