

PRODUCT INFORMATION

Warning: Undefined variable ShacAttributeValueDescription in CAwwwroottmirror.dlmablo.com\wp-content\plugins\woocommerce-print-products\public\class-woocommerce-print-products-public.php on line 2806 verekitug

Unconjugated CRL2, ILXR, TSLPR Conjugate Synonyms Applications ELISA, Flow Cyt

Recommend Dilutions ELISA 1:5000-10000, Flow Cyt 1:100

Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution. Formulation & Reconstitution

Human IgG1(E356D,M358L) - kappa IgG type

Reactivity Target CRLF2 Uniprot ID

Description Anti-CRLF2(verekitug biosimilar) mAb

Delivery

In Stock
Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized antibodies are shipped at ambient temperature.

Background Research grade biosimilar. Not for use in therapeutic or diagnostic procedures for humans or animals

Usage

Anti-CRLF2(verekitug biosimilar) mAb ELISA

0.2 μg of Human CRLF2, His tagged protein per well




Figure 1. ELISA plate pre-coated by 2 μ g/mL (100 μ L/well) Human CRLF2 Protein, His Tag (PME100213) can bind Anti-CRLF2(verekitug biosimilar) mAb (BME100723) in a linear range of 0.64–80 ng/mL.



Email: info@dimabio.com Website: www.dimabio.com








Figure 2. Flow cytometry analysis with 1 μ g/mL Anti-CRLF2(verekitug biosimilar) mAb (BME100723) on HEK293 cells transfected with Human CRLF2 and IL7RA protein (Blue histogram) or HEK293 transfected with IL7RA protein (Red histogram).

Email: info@dimabio.com Website: www.dimabio.com

