

PRODUCT INFORMATION

Warning: Undefined variable \$hasAttributeValueDescription in C\u00e4wwroot\u00fanirror.dimabio.com\u00e4wp-content\plugins\u00e4woocommerce-printproducts\u00e4public\class-woocommerce-print-products-public.php on line 2806
280 Clone ID

CXCL10

C7; IFI10; INP10; IP-10; crg-2; mob-1; SCYB10; gIP-10

Host Species

Anti-CXCL10 antibody(2A5), Rabbit mAb Description

Delivery In Stock Uniprot ID P02778 lgG type Rabbit IgG Clonality Monoclonal Reactivity Human ELISA Applicati

Recommen Dilutions ELISA 1/5000-10000

Purification Purified from cell culture supernatant by affinity chromatography

Lyophilized from sterile PBS, pH 7.4. Normally 5 % – 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis fo specific instructions of reconstitution. Formulation & Reconstitution

Storage & Shipping

Scre at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at mishient temperature.

This antimicrobial gene encodes a chemokine of the CXE subfamily and ligand for the receptor CXCR3. Binding of this protein to CXCR3 results in pleiotropic effects, including stimulation of monocytes, natural killer and T-cell migration, and modulation of adhesion molecule expression. This gene may also be a key regulator of the 'cytokine's torm' immune response to SARS-CoV2 infection. [provided by RefSeq. Sep 2020]

Usage Research use only

Anti-CXCL10 (2A5) mAb ELISA 0.1µg of Human CXCL10, hFc tagged protein per well




Figure 1. ELISA plate pre-coated by 1 μ g/ml (100 μ l/well) Human CXCL10 Protein can bind Rabbit anti-CXCL10 monoclonal antibody(clone: 2A5) in a linear range of 5-100 ng/ml.

Email: info@dimabio.com Website: www.dimabio.com



