

## PRODUCT INFORMATION

<b>Common Name</b>	BMS-936564, MDX-1338
<b>Synonyms</b>	CD184;D2S201E;FB22;HM89;HSY3RR;LAP-3;LAP3;LCR1;LESTR;NPY3R;NPYR;NPYRL;NPYY3R;WHIM;WHIMS
<b>Applications</b>	Flow Cyt
<b>Recommended Dilutions</b>	Flow Cyt 1:100
<b>Formulation &amp; Reconstitution</b>	Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.
<b>Host Species</b>	Homo sapiens
<b>IgG type</b>	IgG4
<b>Reactivity</b>	Human
<b>Target</b>	CXCR4
<b>Uniprot ID</b>	P61073
<b>Description</b>	Anti-CXCR4(ulocuplumab biosimilar) mAb
<b>Delivery</b>	In Stock
<b>Storage &amp; Shipping</b>	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
<b>Background</b>	Research grade biosimilar. Not for use in therapeutic or diagnostic procedures for humans or animals.
<b>Usage</b>	Research use only
<b>Conjugate</b>	Unconjugated
<b>DIMA Disclaimer</b>	All DIMA recombinant antibodies are genuinely generated by DIMA Biotech. They are all under patent application. Any protein sequencing or reverse engineering attempt is prohibited. We are actively scrutinizing all patent application to ensure no IP infringement.




Figure 1. Flow cytometry analysis with 15 µg/mL Anti-CXCR4 (ulocuplumab biosimilar) mAb (BME100101) on Expi293 cells transfected with Human CXCR4 protein (Blue histogram) or Expi293 transfected with irrelevant protein (Red histogram).






Figure 2. Flow cytometry analysis of antigen binding of anti-human CXCR4 mAb(BME100101).

(A) BME100101 does not bind to hepG2 cells that do not express CXCR4.

(B) A clear peak shift of BME100101 was seen compared to the control when incubated with CXCR4-expressing HeLa cells, indicating strong binding of BME100101 to CXCR4. Antibodies were incubated at 5 µg/mL.

DIMABIO CONFIDENTIAL

