

## **PRODUCT INFORMATION**

Clone ID 1A5 Dxd **Target** N.A. **Synonyms Host Species** Rabbit

Description Anti-Dxd antibody(1A5); Rabbit mAb

**Delivery** In Stock **Uniprot ID** N.A.

IgG type Rabbit IgG Clonality Monoclonal

N.A. Reactivity **Applications ELISA** 

Recommended **Dilutions** 

ELISA 1:5000-10000

Purified from cell culture supernatant by affinity **Purification** 

chromatography

Formulation & Reconstitution

**Background** 

Storage & Shipping

Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.

Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not

intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient

temperature.

Dxd is a potent anti-cancer drug, acting as an alkaloid camptothecin analog and a DNA topoisomerase I inhibitor. It forms a stable complex with topoisomerase I and DNA, impeding

DNA replication and inducing cancer cell death. Dxd is utilized as a payload in specific antibody-drug conjugates (ADCs), with examples such as trastuzumab deruxtecan targeting HER2 and

patritumab deruxtecan targeting HER3.

Usage Research use only

Conjugate Unconjugated

All DIMA recombinant antibodies are genuinely generated by DIMA Biotech. They are all under

Email: info@dimabio.com Website: www.dimabio.com

patent application. Any protein sequencing or **DIMA Disclaimer** reverse engineering attempt is prohibited. We are

actively scrutinizing all patent application to

ensure no IP infringement.



Address: Wuhan institute of Biotechnology B7, Biolake No.666 Gaoxin Road, Wuhan, Hubei, China Telephone: +1 2409940618(USA) /+86-18062749453(China) /+86-400-006-0995(China)

Cat. No. DME101025



## ELISA assay to evaluate Anti-Dxd Antibody 0.2µg Human IgG-Dxd per well




Figure 1. Elisa plates were pre-coated with IgG-Dxd (0.2 $\mu$ g/per well). Serial diluted anti-Dxd monoclonal antibody (DME101025) solutions were added, washed, and incubated with secondary antibody before Elisa reading. From above data, the EC50 for anti-Dxd monoclonal antibody binding with IgG-Dxd is 3.447ng/ml.

Email: info@dimabio.com Website: www.dimabio.com

