

PRODUCT INFORMATION

Clone ID 1E6 Dxd **Target** N.A. **Synonyms Host Species** Rabbit

Description Anti-Dxd antibody(1E6); Rabbit mAb

Delivery In Stock **Uniprot ID** N.A.

IgG type Rabbit IgG Clonality Monoclonal

N.A. Reactivity **Applications ELISA**

Recommended **Dilutions**

ELISA 1:5000-10000

Purified from cell culture supernatant by affinity **Purification**

chromatography

Formulation & Reconstitution

Background

Storage & Shipping

Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.

Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not

intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient

temperature.

Dxd is a potent anti-cancer drug, acting as an alkaloid camptothecin analog and a DNA topoisomerase I inhibitor. It forms a stable complex with topoisomerase I and DNA, impeding

DNA replication and inducing cancer cell death. Dxd is utilized as a payload in specific antibody-drug conjugates (ADCs), with examples such as trastuzumab deruxtecan targeting HER2 and

patritumab deruxtecan targeting HER3.

Usage Research use only

Conjugate Unconjugated

> All DIMA recombinant antibodies are genuinely generated by DIMA Biotech. They are all under patent application. Any protein sequencing or

> > Email: info@dimabio.com Website: www.dimabio.com

DIMA Disclaimer reverse engineering attempt is prohibited. We are

actively scrutinizing all patent application to

ensure no IP infringement.



Address: Wuhan institute of Biotechnology B7, Biolake No.666 Gaoxin Road, Wuhan, Hubei, China Telephone: +1 2409940618(USA) /+86-18062749453(China) /+86-400-006-0995(China)



ELISA assay to evaluate Anti-Dxd Antibody 0.2µg Human IgG-Dxd per well




Figure 1. Elisa plates were pre-coated with IgG-Dxd (0.2 μ g/per well). Serial diluted anti-Dxd monoclonal antibody (DME101027) solutions were added, washed, and incubated with secondary antibody before Elisa reading. From above data, the EC50 for anti-Dxd monoclonal antibody binding with IgG-Dxd is 68.68ng/ml.

Email: info@dimabio.com Website: www.dimabio.com

