

## PRODUCT INFORMATION

Warning: Undefined variable ShasAttributeValueDescription in C:\www.root\mirror.dimablo.com\wp-content\plugins\woocommerce-print-products\public.class-woocommerce-print-products-public.php on line 2806 bMC472 Clone ID

EK8; HEK8; SEK; TYRO1 Synonyme

Host Species Rabbit

Anti-EPHA4 antibody(DMC472); IgG1 Chimeric mAb Description

Delivery In Stock Uniprot ID P54764

Rabbit/Human Fc chimeric IgG1 lgG type

Clonality Monoclonal Reactivity Human Application Flow Cyt

Recommend Dilutions Flow Cyt 1:100

Purification Purified from cell culture supernatant by affinity chromatography

Formulation & Reconstitution Lyophilized from sterile PBS, pH 7.4. Normally 5 % – 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.

Storage & Shipping

specific instructions of reconstitution.

Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.

This gene belongs to the ephrin receptor subfamily of the protein-tryonsine kinase family, EPH and EPH-related receptors have been implicated in mediating developmental events; particularly in the nervous system. Receptors in the EPH subfamily typically have a single kinase domain and an extracellular region containing a Cys-rich domain and 2 fibronectin type III repeats. The ephrin receptors are dividen to 2 groups based on the similarity of their extracellular domain sequences and their affinities for binding ephrin-A and ephrin-B ligands. Multiple transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq; jan 2015]

Research use only Background

Usage Coniugate

All DIMA recombinant antibodies are genuinely generated by DIMA Biotech. They are all under patent application. Any protein sequencing or reverse engineering attempt is prohibited. We are actively scrutinizing all patent application to ensure no IP infringement. DIMA Disclaime




Figure 1. Flow cytometry analysis with Anti-EPHA4 (DMC472) on HEK293 cells transfected with human EPHA4 (Blue histogram) or HEK293 transfected with irrelevant protein (Red histogram).

Address: Wuhan institute of Biotechnology B7, Biolake No.666 Gaoxin Road, Wuhan, Hubei, China Telephone: +1 2409940618(USA) /+86-18062749453(China) /+86-400-006-0995(China)

Email: info@dimabio.com Website: www.dimabio.com



Cat. No. DMC100472






Figure 2. Flow cytometry analysis of antigen binding of anti-human EPHA4 mAb(DMC100472). (A) DMC100472 does not bind to CHO-S cells that do not express EPHA4. (B) A clear peak shift of DMC100472 was seen compared to the control when incubated with EPHA4-expressing MCF-7 cells, indicating strong binding of DMC100472 to EPHA4. Antibodies were incubated at 5  $\mu$ g/mL.





