

PRODUCT INFORMATION

Warning . Undefined variable \$hasAttributeValueDescription in Calwawroottmirror.dimablo.com/wp-content\plugins/woocommerce-print-products-public.php on line 2806 [5]/21

Conjugate Synonyms Applications ELISA, Flow Cyt

Recommend Dilutions ELISA 1:5000-10000, Flow Cyt 1:100

Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution. Formulation & Reconstitution

Human IgG1(L234A,L235A) - Kappa IgG type

Reactivity Target FCGR3A Uniprot ID

Anti-FCGR3A(AFM13 without CD30 biosimilar) mAb Description

Delivery


Background

In Stock
Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized antibodies are shipped at ambient temperature.

Usage

Anti-FCGR3A(AFM13 without CD30 biosimilar) mAb ELISA

0.2 μg of Human FCGR3A, His tagged protein per well



Anti-FCGR3A (AFM13 without CD30 biosimilar) antibody(ng/ml)

Figure 1. ELISA plate pre-coated by 2 μ g/mL (100 μ L/well) Human FCGR3A Protein (F176V), His Tag (PME100069) can bind Anti-FCGR3A(AFM13 without CD30 biosimilar) mAb (BME100267) in a linear range of 0.64–80 ng/mL.

Email: info@dimabio.com Website: www.dimabio.com










Figure 2. Flow cytometry analysis with 1 μ g/mL Anti-FCGR3A(AFM13 without CD30 biosimilar) mAb (BME100267) on HEK293 cells transfected with Human FCGR3A protein (Blue histogram) or HEK293 transfected with irrelevant protein (Red histogram).

Email: info@dimabio.com Website: www.dimabio.com

