

PRODUCT INFORMATION

Clone ID 18A3 FOLR1 **Target**

Synonyms FBP;FOLR;FRalpha

Host Species Rabbit

Description Anti-FOLR1 antibody(18A3), IgG1 Chimeric mAb

Delivery In Stock **Uniprot ID** P15328

IgG type Rabbit/Human Fc chimeric IgG1

Clonality Monoclonal Reactivity Human **Applications** WB

Recommended

Storage & Shipping

WB 1:1000 **Dilutions**

Purified from cell culture supernatant by affinity **Purification**

chromatography

Lyophilized from sterile PBS, pH 7.4. Normally 5 % Formulation & - 8% trehalose is added as protectants before Reconstitution

lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution. Store at -20°C to -80°C for 12 months in

lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing).

Lyophilized proteins are shipped at ambient

temperature.

The protein encoded by this gene is a member of the folate receptor family. Members of this gene family bind folic acid and its reduced derivatives, and transport 5-methyltetrahydrofolate into cells. This gene product is a secreted protein that either

anchors to membranes via a glycosyl-

phosphatidylinositol linkage or exists in a soluble **Background**

form. Mutations in this gene have been associated with neurodegeneration due to cerebral folate transport deficiency. Due to the presence of two promoters, multiple transcription start sites, and alternative splicing, multiple transcript variants encoding the same protein

have been found for this gene.

Usage Research use only

Conjugate Unconjugated

All DIMA recombinant antibodies are genuinely generated by DIMA Biotech. They are all under

patent application. Any protein sequencing or **DIMA Disclaimer** reverse engineering attempt is prohibited. We are

actively scrutinizing all patent application to

ensure no IP infringement.



Email: info@dimabio.com Website: www.dimabio.com








Figure 1.Anti-FOLR1 antibody (SKU# DMC100585) at 1/1000 dilution

Lane: HeLa (human cervical adenocarcinoma epithelial cell), whole cell lysate

Secondary: Goat Anti-Rabbit IgG H&L (HRP) at 1/5000 dilution

band size: 38 kDa




Figure 2. Flow cytometry analysis of antigen binding of anti-human FOLR1 mAb(DMC100585). (A) DMC100585 does not bind to K562 cells that do not express FOLR1. (B) A clear peak shift of DMC100585 was seen compared to the control when incubated with FOLR1-expressing Hela cells, indicating strong binding of DMC100585 to FOLR1. Antibodies were incubated at 5 μ g/mL.



Email: info@dimabio.com Website: www.dimabio.com

