

## PRODUCT INFORMATION

<b>Clone ID</b>	<b>Warning:</b> Undefined variable \$hasAttributeValueDescription in C:\wwwroot\mirror.dimabio.com\wp-content\plugins\woocommerce-print-products\public\class-woocommerce-print-products-public.php on line 2806 12H7
<b>Target</b>	G4S linker
<b>Synonyms</b>	GGGG5
<b>Host Species</b>	Rabbit
<b>Description</b>	Anti-(G4S)4 antibody(12H7), Rabbit mAb
<b>Delivery</b>	In Stock
<b>Uniprot ID</b>	N/A
<b>IgG type</b>	Rabbit IgG
<b>Clonality</b>	Monoclonal
<b>Reactivity</b>	N/A
<b>Applications</b>	ELISA
<b>Recommended Dilutions</b>	ELISA 1/5000-10000
<b>Purification</b>	Purified from cell culture supernatant by affinity chromatography
<b>Formulation &amp; Reconstitution</b>	Lyophilized from sterile PBS, pH 7.4. Normally 5% - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.
<b>Yefei Storage</b>	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
<b>Background</b>	The poly-Glycine-Serine (G4S) linker is a type of flexible, unstructured synthetic peptide linker sequence often leveraged to connect antibody fragments (scFvs) and fusion proteins. The linker itself consists of a core pentapeptide sequence, Gly-Gly-Gly-Ser, that is repeated and commonly found as either a 15-mer (G4S)3 or 20-mer (G4S)4 within scFv-based CARs and scFv fragments. The linker sequence length plays a role in controlling scFv stability and the noncovalent association between the VH and VL domains.
<b>Usage</b>	Research use only
<b>Conjugate</b>	Unconjugated




Figure 1. ELISA plate pre-coated by 1 µg/ml (100 µl/well) Human (G4S)4, KLH tagged protein can bind Rabbit anti-(G4S)4 monoclonal antibody(clone: 12H7) in a linear range of 10-100 ng/ml.

