

## **PRODUCT INFORMATION**

iable \$hasAttributeValueDescription in C:\wwwroot\mirror.dimabio.com\wp-content\plugins\woocommerce-print-woocommerce-print-products-public.php on line 2806

Conjugate

Unconjugated
GM-CSF-R-alpha; GMCSFR-alpha; GMR-alpha; CDw116; CD116; CSF2RA; CSF2R; CSF2RY Synonyms

Applications Flow Cyt Recommend Dilutions Flow Cyt 1:100

Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution. Formulation & Reconstitution

Human IgG4(S228P) - Kappa IgG type

Reactivity Target GM-CSFR Uniprot ID

Anti-GM-CSFR(mavrilimumab biosimilar) mAb Description

Delivery

Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized antibodies are shipped at ambient temperature.

Background Research grade biosimilar. Not for use in therapeutic or diagnostic procedures for humans or animals

Usage




Figure 1. Flow cytometry analysis with  $15\mu g/mL$  Anti-GM-CSFR(mavrilimumab biosimilar) mAb (BME100213) on HEK293 cells transfected with Human GM-CSFR protein (Blue histogram) or HEK293 transfected with irrelevant protein (Red histogram).

Address: Wuhan institute of Biotechnology B7, Biolake No.666 Gaoxin Road, Wuhan, Hubei, China Telephone: +1 2409940618(USA) /+86-18062749453(China) /+86-400-006-0995(China)

Email: info@dimabio.com Website: www.dimabio.com

