

PRODUCT INFORMATION

Warning . Undefined variable \$hasAttributeValueDescription in Calwawroottmirror.dimabio.com/wp-content\piugins/woocommerce-print-mcARH109*
MCARH109*

WARH109*

WARH10

Conjugate Synonyms Applications ELISA, Flow Cyt

Recommend Dilutions ELISA 1:5000-10000, Flow Cyt 1:100

Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution. Formulation & Reconstitution

IgG type Human IgG1 - kappa Reactivity

GPRC5D Uniprot ID Anti-GPRC5D(MCARH109 biosimilar) mAb Description

Delivery

Target

Background

Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized antibodies are shipped at ambient temperature.

Research grade biosimilar. Not for use in therapeutic or diagnostic procedures for humans or animals

Usage

Anti-GPRC5D(MCARH109 biosimilar) mAb ELISA

 $0.2~\mu g$ of Human GPRC5D, mFc tagged protein per well




Figure 1. ELISA plate pre-coated by 5 μ g/mL (100 μ L/well) Human GPRC5D Protein, mFc Tag (PME100559) can bind Anti-GPRC5D(MCARH109 biosimilar) mAb (BME100265) in a linear range of 16–2000 ng/mL.

Email: info@dimabio.com Website: www.dimabio.com








Figure 2. Flow cytometry analysis with 1 μ g/mL Anti-GPRC5D(MCARH109 biosimilar) mAb (BME100265) on HEK293 cells transfected with Human GPRC5D protein (Blue histogram) or HEK293 transfected with irrelevant protein (Red histogram).

Email: info@dimabio.com Website: www.dimabio.com

