

## **PRODUCT INFORMATION**

Clone ID 6E3

**GUCY2C Target** 

**Synonyms** DIAR6;GC-C;GUC2C;MECIL;MUCIL;STAR

**Host Species** Rabbit

Description Anti-GUCY2C antibody(6E3), IgG1 Chimeric mAb

**Delivery** In Stock **Uniprot ID** P25092

IgG type Rabbit/Human Fc chimeric IgG1

Clonality Monoclonal Reactivity Human **Applications** WB

Recommended

**Dilutions** 

WB 1:1000

Purified from cell culture supernatant by affinity **Purification** 

chromatography

Formulation & Reconstitution

**Background** 

Storage & Shipping

Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.

Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store

at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient

temperature.

This gene encodes a transmembrane protein that functions as a receptor for endogenous peptides guanylin and uroguanylin, and the heat-stable E. coli enterotoxin. The encoded protein activates the cystic fibrosis transmembrane conductance regulator. Mutations in this gene are associated

with familial diarrhea (autosomal dominant) and meconium ileus (autosomal recessive). [provided by RefSeq, Nov 2016]

Usage Research use only

Conjugate Unconjugated

> All DIMA recombinant antibodies are genuinely generated by DIMA Biotech. They are all under patent application. Any protein sequencing or

> > Email: info@dimabio.com Website: www.dimabio.com

**DIMA Disclaimer** reverse engineering attempt is prohibited. We are

actively scrutinizing all patent application to

ensure no IP infringement.








Figure 1.Anti-GUCY2C antibody (SKU# DMC100508) at 1/1000 dilution

Lane: HT55, whole cell lysate

Secondary: Goat Anti-Rabbit IgG H&L (HRP) at 1/5000 dilution

Predicted band size: 123 kDa Observed band size: 150 kDa



