

## **PRODUCT INFORMATION**

Clone ID 55A1
Target HER3

Synonyms ERBB3; FERLK; LCCS2; VSCN1; ErbB-3; c-erbB3; erbB3-S; MDA-BF-1; c-erbB-3; p180-ErbB3; p45-

sErbB3; p85-sErbB3

Host Species Rabbit

**Description** Anti-Her3 antibody(55A1), Rabbit mAb

Delivery In Stock
Uniprot ID P21860
IgG type Rabbit IgG
Clonality Monoclonal
Reactivity Human

Applications IHC; Flow Cyt

Recommended Dilutions

Storage & Shipping

Background

IHC 1:100; Flow Cyt 1:100

**Purification**Purified from cell culture supernatant by affinity

chromatography

Lyophilized from sterile PBS, pH 7.4. Normally 5 %
- 8% trehalose is added as protectants before
lyophilization. Please see Certificate of Analysis

for specific instructions of reconstitution. Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store

at -80°C (Avoid repeated freezing and thawing).
Lyophilized proteins are shipped at ambient

témperature.

This gene encodes a member of the epidermal growth factor receptor (EGFR) family of receptor tyrosine kinases. This membrane-bound protein has a neuregulin binding domain but not an active kinase domain. It therefore can bind this ligand but not convey the signal into the cell through protein phosphorylation. However, it does form heterodimers with other EGF receptor family members which do have kinase activity. Heterodimerization leads to the activation of pathways which lead to cell proliferation or differentiation. Amplification of this gene and/or overexpression of its protein have been reported

overexpression of its protein have been reported in numerous cancers, including prostate, bladder, and breast tumors. Alternate transcriptional splice variants encoding different isoforms have been


characterized. One isoform lacks the

intermembrane region and is secreted outside the cell. This form acts to modulate the activity of the membrane-bound form. Additional splice variants have also been reported, but they have not been thoroughly characterized. [provided by RefSeq, Jul

Email: info@dimabio.com Website: www.dimabio.com

2008]

Usage Research use only
Conjugate Unconjugated





Cat. No. DME100580



**DIMA Disclaimer** 

All DIMA recombinant antibodies are genuinely generated by DIMA Biotech. They are all under patent application. Any protein sequencing or reverse engineering attempt is prohibited. We are actively scrutinizing all patent application to ensure no IP infringement.




Figure 1. Flow cytometry analysis of antigen binding of anti-human Her3 mAb(DME100580).

(A) DME100580 does not bind to 293T cells that do not express Her3
(B) A clear peak shift of DME100580 was seen compared to the control when incubated with Her3-expressing Huh7 cells, indicating strong binding of DME100580 to Her3.

Email: info@dimabio.com Website: www.dimabio.com

Antibodies were incubated at 10 µg/mL.

