

PRODUCT INFORMATION

Warning: Undefined variable ShasAttributeValueDescription in C:\www.root\mirror.dimablo.com\wp-content\plugins\woocommerce-print-products\public\class-woocommerce-print-products-public.php on line 2806 R0485896-000.teprotumumab-trbw

Conjugate Unconjugated CD221, IGFIR, IGFR, JTK13 Synonyms Applications Flow Cyt Recommende Dilutions Flow Cyt 1:100

Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions. Formulation & Reconstitution

Human IgG1(E356D,M358L) - kappa IgG type

Reactivity Target IGF-1R Uniprot ID

Description Anti-IGF-1R(teprotumumab biosimilar) mAb

Delivery Storage & Sh

Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized antibodies are shipped at ambient temperature.

Research grade biosimilar. Not for use in therapeutic or diagnostic procedures for humans or animals

Background Usage




Figure 1. IGF-1R protein is highly expressed on the surface of HEK293 cell membrane. Flow cytometry analysis with $1\mu g/mL$ Anti-IGF-1R(teprotumumab biosimilar) mAb (BME100174) on HEK293 cells transfected with human IGF-1R (Blue histogram) or HEK293 transfected with irrelevant protein (Red histogram), and Isotype antibody on HEK293 transfected with irrelevant protein (Orange histogram).

Email: info@dimabio.com Website: www.dimabio.com

Address: Wuhan institute of Biotechnology B7, Biolake No.666 Gaoxin Road, Wuhan, Hubei, China Telephone: +1 2409940618(USA) /+86-18062749453(China) /+86-400-006-0995(China)








Figure 2. Flow cytometry analysis of antigen binding of anti-human IGF-1R mAb(BME100174). (A) BME100174 does not bind to Jurkat cells that do not express IGF-1R. (B) A clear peak shift of BME100174 was seen compared to the control when incubated with IGF-1R-expressing Hela cells, indicating strong binding of BME100174 to IGF-1R. Antibodies were incubated at $5 \mu g/mL$.




Figure 3. Flow cytometry analysis of antigen binding of anti-human IGF-1R mAb(BME100174). (A) BME100174 does not bind to Jurkat cells that do not express IGF-1R. (B) A clear peak shift of BME100174 was seen compared to the control when incubated with IGF-1R-expressing MCF-7 cells, indicating strong binding of BME100174 to IGF-1R. Antibodies were incubated at 5 μ g/mL.

Email: info@dimabio.com Website: www.dimabio.com

