

PRODUCT INFORMATION

Warning: Undefined variable ShasAttributeValueDescription in C:\www.root\mirror.dimabio.com\wp-content\plugins\woocommerce-print-products\public\class-woocommerce-print-products-public.php on line 2806 bM125 Clone ID

II 17RΔ

CD217;CDw217;IL-17RA;IL17R;CANDF5;hIL-17R

Host Species Rabbit

Anti-IL17RA antibody(DM125); Rabbit mAb Description In Stock

Delivery Q96F46 Uniprot ID IgG type Rabbit IgG Clonality Monoclonal Reactivity Human Flow Cyt Application Recommend Dilutions Flow Cyt 1:100

Purification Purified from cell culture supernatant by affinity chromatography

Formulation & Reconstitution Lyophilized from sterile PBS, pH 7.4. Normally 5 % – 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.

Storage & Shipping

specific instructions of reconstitution.

Store at 20°C to 80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.

Interleukin 174 (IL-174) is a priorinflammatory cytokine secreted by activated T-lymphocytes. It is a potent inducer of the maturation of CD34-positive hematopoietic precursors into neutrophils. The transmembrane protein encoded by this gene (interleukin 174 receptor; IL17A) is a ubiquitious type I membrane glycoprotein that binds with low affinity to interleukin 174 and rescreptor play appeal proposed protein and autoimmune diseases such as rheumatoid arthritis. Like other cytokine receptors; this receptor lakely has a multimeric structure. Alternative splicing results in multiple transcript variants encoding different isoforms.

Research use only Background

Usage

Coniugate

All DIMA recombinant antibodies are genuinely generated by DIMA Biotech. They are all under patent application. Any protein sequencing or reverse engineering attempt is prohibited. We are actively scrutinizing all patent application to ensure no IP infringement. DIMA Disclaime




Figure 1. Flow cytometry analysis with Anti-IL-17RA (DM125) on HEK293 cells transfected with human IL17RA (Blue histogram)or HEK293 transfected with irrelevant protein (Red histogram).











Figure 2. Flow cytometry analysis of antigen binding of rabbit anti-human IL17RA mAb(DME100125). (A) DME100125 does not bind to CHO-S cells that do not express IL17RA. (B) A clear peak shift of DME100125 was seen compared to the control when incubated with IL17RA-expressing THP-1 cells, indicating strong binding of DME100125 to IL17RA. Antibodies were incubated at 5 μ g/mL.



Email: info@dimabio.com Website: www.dimabio.com

