

PRODUCT INFORMATION

Warning: Undefined variable ShasAttributeValueDescription in C:\u00e4wroot\u00fcmirror.dimablo.com\u00e4wp-content\plugins\u00e4woocommerce-print-products\u00e4public\u00e7class-woocommerce-print-products-public.php on line 2806 bb/206 Clone ID

II 15RA

CD215 Synonyme Host Species Rabbit

Anti-IL15RA antibody(DM206); Rabbit mAb Description Delivery In Stock

Uniprot ID Q13261 lgG type Rabbit IgG Clonality Monoclonal Reactivity Human Application ELISA; Flow Cyt

Recommend Dilutions ELISA 1:5000-10000; Flow Cyt 1:100

Purification Purified from cell culture supernatant by affinity chromatography

Formulation & Reconstitution Lyophilized from sterile PBS, pH 7.4. Normally 5 % – 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.

Storage & Shipping

specific instructions of reconstitution.

Store at 2-0°C to 80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.

This gene encodes a cytokine receptor that specifically binds interreduciin 15 (ILIS) with high affinity. The receptors of ILIS and IL2 share two subunits; ILIZR beta and IL2R gamma. This forms the basis of many overlapping biological activities of ILIS and IL2. The protein encoded by this gene is the protein encoded by this gene is the protein encoded by the gene is the gene is the protein encoded by the gene is the gene g

Usage Research use only

Inconjugated
All DIMA recombinant antibodies are genuinely generated by DIMA Blotech. They are all under patent application. Any protein sequencing or reverse
engineering attempt is prohibited. We are actively scrutinizing all patent application to ensure no IP infringement. DIMA Disclaimer




Figure 1. Flow cytometry analysis with Anti-IL15RA (DM206) on HEK293 cells transfected with human IL15RA (Blue histogram) or HEK293 transfected with irrelevant protein (Red histogram).

Email: info@dimabio.com Website: www.dimabio.com








Figure 2. Flow cytometry analysis of antigen binding of rabbit anti-human IL15RA mAb(DME100206). (A) DME100206 does not bind to CHO-S cells that do not express IL15RA. (B) A clear peak shift of DME100206 was seen compared to the control when incubated with IL15RA-expressing Siha cells, indicating strong binding of DME100206 to IL15RA. Antibodies were incubated at 5 μ g/mL.



