

PRODUCT INFORMATION

Clone ID 1G1 IL18BP **Target**

Synonyms FVH:IL18BPa

Host Species Rabbit

Description Anti-IL18BP antibody(1G1), IgG1 Chimeric mAb

Delivery In Stock **Uniprot ID** 095998

IgG type Rabbit/Human Fc chimeric IgG1

Clonality Monoclonal Reactivity Human **Applications ELISA**

Recommended **Dilutions**

Background

ELISA 1/5000-10000

Purified from cell culture supernatant by affinity **Purification**

chromatography

Lyophilized from sterile PBS, pH 7.4. Normally 5 % Formulation & - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis Reconstitution

for specific instructions of reconstitution. Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not

intended for use within a month, aliquot and store Storage & Shipping at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient

temperature.

The protein encoded by this gene functions as an inhibitor of the proinflammatory cytokine, IL18. It binds IL18, prevents the binding of IL18 to its receptor, and thus inhibits IL18-induced IFNgamma production, resulting in reduced T-helper type 1 immune responses. This protein is

constitutively expressed and secreted in mononuclear cells. Elevated level of this protein is detected in the intestinal tissues of patients with Crohn's disease. Alternatively spliced transcript variants encoding different isoforms have been described for this gene. [provided by RefSeq, Feb 2011]

Usage Research use only Conjugate Unconjugated

> All DIMA recombinant antibodies are genuinely generated by DIMA Biotech. They are all under patent application. Any protein sequencing or

> > Email: info@dimabio.com Website: www.dimabio.com

DIMA Disclaimer reverse engineering attempt is prohibited. We are actively scrutinizing all patent application to

ensure no IP infringement.

Address: Wuhan institute of Biotechnology B7, Biolake No.666 Gaoxin Road, Wuhan, Hubei, China Telephone: +1 2409940618(USA) /+86-18062749453(China)

/+86-400-006-0995(China)





Anti-IL18BP (1G1) mAb ELISA 0.1µg of IL18BP, hFc tagged protein per well




Figure 1. ELISA plate pre-coated by 1 μ g/ml (100 μ l/well) Human IL18BP, hFc tagged protein can bind Rabbit anti-IL18BP monoclonal antibody(clone: 1G1) in a linear range of 5-100 ng/ml.

