

PRODUCT INFORMATION

riable \$hasAttributeValueDescription in C:\wwwroot\mirror.dimabio.com\wp-content\plugins\woocommerce-print-woocommerce-print-products-public.php on line 2806 Clone ID

II 31RΔ

CRL; GPL; CRL3; GLMR; GLM-R; PLCA2; hGLM-R; IL-31RA; PRO21384; zcytoR17

Host Species

Anti-IL31RA antibody(2A6), IgG1 Chimeric mAb Description

Delivery In Stock Uniprot ID Q8NI17

Rabbit/Human Fc chimeric IgG1 lgG type

Clonality Monoclonal Reactivity Human ELISA Applicati

Recommen Dilutions ELISA 1/5000-10000

Purification Purified from cell culture supernatant by affinity chromatography

Formulation & Reconstitution Lyophilized from sterile PBS, pH 7.4. Normally 5 % – 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.

Storage & Shipping

Background

specinic instructions of reconstitution.

Store at -20°C to -80°C for 12 months in Jophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.

The protein encoded by this gene belongs to the type I cytokine receptor family. This receptor, with homology to gp130, is expressed on monocytes, and is involved in II-31 signaling via activation of STAT-3 and STAT-5. It functions either as a monomer, or as part of a receptor complex with oncostatin M receptor (OSMR). Several alternatively spliced transcript variants encoding different isoforms have been noted for this gene_Iprovided by RefSeq, Jun 2011]

Usage Conjugate Unconjugated

Anti-IL31RA (2A6) mAb ELISA 0.1µg of IL31RA, mFc tagged protein per well




Figure 1. ELISA plate pre-coated by 1 μ g/ml (100 μ l/well) Human IL31RA, mFc tagged protein can bind Rabbit anti-IL31RA monoclonal antibody(clone: 2A6) in a linear range of 1-10 ng/ml.



Email: info@dimabio.com Website: www.dimabio.com

