

## PRODUCT INFORMATION

Warning: Undefined variable ShasAttributeValueDescription in C:\www.root\mirror.dimablo.com\wp-content\plugins\woocommerce-print-products\public.class-woocommerce-print-products-public.php on line 2806 bMC284 Clone ID

CD124; IL-4RA; IL4RA Synonyme

Host Species Rabbit

Anti-IL4RA antibody(DMC284); IgG1 Chimeric mAb Description

Delivery In Stock

Uniprot ID P24394 Rabbit/Human Fc chimeric IgG1 lgG type

Clonality Monoclonal Reactivity Human Application Flow Cyt Recommend Dilutions Flow Cyt 1:100

Purification Purified from cell culture supernatant by affinity chromatography

Formulation & Reconstitution Lyophilized from sterile PBS, pH 7.4. Normally 5 % – 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.

Storage & Shipping

specific instructions of reconstitution.

Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.

This gene encodes the alight achian of the interfeukin-4 receptor, a type I transmembrane protein that can bind interieukin and interieukin 13 to This gene encodes the alight achian of the interfeukin-4 receptor, a type I transmembrane protein that can bind interieukin and interieukin 13 to This gene encodes the alight and the interiously protein can be produced by proteolysis of the membrane-bound protein; and this soluble form can inhibit IL4-mediated cell proliferation and IL5 upregulation by T-cells. Allelic variations in this gene have been associated with actory; a condition that can manifest itself as allergic inhibits; sinsubus; asthma; or eczema. Polymorphisms in this gene are also associated with resistance to human immunodeficiency virus type-1 infection. Alternate splicing results in multiple transcript variants.

Usage Research use only

Inconjugated
All DIMA recombinant antibodies are genuinely generated by DIMA Blotech. They are all under patent application. Any protein sequencing or reverse
engineering attempt is prohibited. We are actively scrutinizing all patent application to ensure no IP infringement. DIMA Disclaimer




Figure 1. Flow cytometry analysis with Anti-IL4RA (DMC284) on HEK293 cells transfected with human IL4RA (Blue histogram) or HEK293 transfected with irrelevant protein (Red histogram).

Address: Wuhan institute of Biotechnology B7, Biolake No.666 Gaoxin Road, Wuhan, Hubei, China Telephone: +1 2409940618(USA) /+86-18062749453(China) /+86-400-006-0995(China)

Email: info@dimabio.com Website: www.dimabio.com

