

## **PRODUCT INFORMATION**

Warning: Undefined variable ShasAttributeValueDescription in C:\u00e4wroot\u00e4mirror.dimablo.com\u00e4wp-content\plugins\u00e4woocommerce-print-products\u00e4public\u00e7pablic\u00e4cass-woocommerce-print-products-public.php on line 2806 bMC274 Clone ID

Synonyme

Host Species Rabbit

Anti-IL5 antibody(DMC274); IgG1 Chimeric mAb Description

Delivery In Stock Uniprot ID P05113

Rabbit/Human Fc chimeric IgG1 lgG type

Clonality Monoclonal Reactivity Human Flow Cyt Application Recommend Dilutions Flow Cyt 1:100

Purification Purified from cell culture supernatant by affinity chromatography

Formulation & Reconstitution Lyophilized from sterile PBS, pH 7.4. Normally 5 % – 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.

Storage & Shipping

Specific instructions of reconstitution.

Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at antiheline temperature.

This gene encodes a cytokine that acts as a growth and differentiation factor for both B cells and easinophils. The encoded cytokine plays a major role pathogenesis of eosinophil-dependent inflammatory diseases. This cytokine functions by binding to its receiption, which is a heterodimer; whose beat subunit is shared with the receiptors for interleukine 3 (II.3) and colony stimulating factor 2 (CSF-SM-CSF). This gene is located on chromosome 5 with a cytokine gene cluster which includes interleukin 4 (II.4); interleukin 13 (II.3); and CSF2. This gene; II.4; and II.13 may be regulated coordinately by long-grange regulatory elements spread over 120 citiobases on chromosome 65 with

Usage Research use only

Inconjugated
All DIMA recombinant antibodies are genuinely generated by DIMA Blotech. They are all under patent application. Any protein sequencing or reverse
engineering attempt is prohibited. We are actively scrutinizing all patent application to ensure no IP infringement. DIMA Disclaimer




Figure 1. Flow cytometry analysis with Anti-IL5 (DMC274) on HEK293 cells transfected with human IL5 (Blue histogram) or HEK293 transfected with irrelevant protein (Red histogram).

Address: Wuhan institute of Biotechnology B7, Biolake No.666 Gaoxin Road, Wuhan, Hubei, China Telephone: +1 2409940618(USA) /+86-18062749453(China) /+86-400-006-0995(China)

Email: info@dimabio.com Website: www.dimabio.com

