

PRODUCT INFORMATION

riable \$hasAttributeValueDescription in C:\wwwroot\mirror.dimabio.com\wp-content\plugins\woocommerce-print-woocommerce-print-products-public.php on line 2806

Conjugate

Unconjugated Activin beta-A chain INHBA, Activin A Synonyms

Applications ELISA

Recommend Dilutions ELISA 1:5000-10000

Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution. Formulation & Reconstitution

IgG type Human IgG4 - Kappa Reactivity Target INHBA Uniprot ID

Description Anti-INHBA(garetosmab biosimilar) mAb

Delivery

Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized antibodies are shipped at ambient temperature.

Background Research grade biosimilar. Not for use in therapeutic or diagnostic procedures for humans or animals

Usage

Anti-INHBA(garetosmab biosimilar) mAb ELISA

0.2 μg of Human Activin A protein per well




Figure 1. ELISA plate pre-coated by 2 μ g/mL (100 μ L/well) Human Activin A Protein can bind Anti-INHBA(garetosmab biosimilar) mAb (BME100519) in a linear range of 2.29–185.19 ng/mL.



Email: info@dimabio.com Website: www.dimabio.com

