

PRODUCT INFORMATION

mon Name Warning: Undefined variable \$hasAttributeValueDescription in C:\www.root\mirror.dimabio.com\wp-content\plugins\woocommerce-print-products-public.class-woocommerce-print-products-public.php on line 2806 A8T-700.

Conjugate Unconjugated
Synonyms MET
Applications ELISA, Flow Cyt

Recommended Dilutions ELISA 1:5000-10000, Flow Cyt 1:100

Formulation & Reconstitution Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.

specific instructions of i

IgG type Human IgG1(K214R,Del-K222,T223C,Del-T225) - Kappa

 Reactivity
 Human

 Target
 MET

 Uniprot ID
 P08581

Description Anti-MET(telisotuzumab biosimilar) mAb

Delivery In Stoc

Background

ping
Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized antibodies are shipped at ambient temperature.

Research grade biosimilar. Not for use in therapeutic or diagnostic procedures for humans or animals

Usage Research use onl

Anti-MET(telisotuzumab biosimilar) mAb ELISA

0.2 μg of Human MET, His tagged protein per well




Figure 1. ELISA plate pre-coated by 2 μ g/mL (100 μ L/well) Human MET Protein, His Tag (PME101405) can bind Anti-MET(telisotuzumab biosimilar) mAb (BME100261) in a linear range of 0.13–16 ng/mL.

Email: info@dimabio.com Website: www.dimabio.com

Address: Wuhan institute of Biotechnology B7, Biolake No.666 Gaoxin Road, Wuhan, Hubei, China Telephone: +1 2409940618(USA) /+86-18062749453(China) /+86-400-006-0995(China)






Figure 2. Flow cytometry analysis with 1 μ g/mL Anti-MET(telisotuzumab biosimilar) mAb (BME100261) (Blue histogram) or isotype control mAb (Red histogram) on HEK293 cells.

Email: info@dimabio.com Website: www.dimabio.com

