

## **PRODUCT INFORMATION**

Warning: Undefined variable ShasAttributeValueDescription in C:\www.root\mirror.dimablo.com\wp-content\plugins\woocommerce-print-products-public.php on line 2806
PABODI[PABOD1-ADC][MUCI-1H7, Unconjugated mab

Conjugate Unconjugated MUC-1; CD227; PUM Synonyms Applications ELISA, Flow Cyt

Recommend Dilutions ELISA 1:5000-10000, Flow Cyt 1:100

Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution. Formulation & Reconstitution

Human IgG1(E356D,M358L) - kappa IgG type

Reactivity Target MUC1 Uniprot ID P15941

Anti-MUC1(PAb-001 biosimilar) mAb Description

Delivery

## Anti-MUC1(PAb-001 biosimilar) mAb ELISA

0.2 μg of Human MUC1, His tagged protein per well




Figure 1. ELISA plate pre-coated by 2  $\mu$ g/mL (100  $\mu$ L/well) Human MUC1 Protein, His Tag (PME100106) can bind Anti-MUC1(PAb-001 biosimilar) mAb (BME100249) in a linear range of 0.64-80 ng/mL.

Email: info@dimabio.com Website: www.dimabio.com

Address: Wuhan institute of Biotechnology B7, Biolake No.666 Gaoxin Road, Wuhan, Hubei, China Telephone: +1 2409940618(USA) /+86-18062749453(China) /+86-400-006-0995(China)








Figure 2. Flow cytometry analysis with  $1\mu g/mL$  Anti-MUC1(PAb-001 biosimilar) mAb (BME100249) on HEK293 cells transfected with Human MUC1 protein (Blue histogram) or HEK293 transfected with irrelevant protein (Red histogram).

