

PRODUCT INFORMATION

Warning: Undefined variable ShasAttributeValueDescription in C:\www.root\mirror.dimablo.com\wp-content\plugins\woocommerce-print-products\public\class-woocommerce-print-products-public.php on line 2806 bMC298 Clone ID

CD159A; NKG2; NKG2A Synonyme

Host Species Rabbit

Anti-NKG2A antibody(DMC298); IgG1 Chimeric mAb Description

Delivery In Stock Uniprot ID P26715

Rabbit/Human Fc chimeric IgG1 lgG type

Clonality Monoclonal Reactivity Human Flow Cyt Applicati Recommend Dilutions Flow Cyt 1:100

Purification Purified from cell culture supernatant by affinity chromatography

Formulation & Reconstitution Lyophilized from sterile PBS, pH 7.4. Normally 5 % – 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.

Storage & Shipping

Specific instructions of reconstitution.

Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at a mibilent temperature.

Astural killer (IMC) cells are lyophocytes that can mediate lysis of certain tumor cells and virue-infected cells without previous activation. They can also Natural killer (IMC) cells are lyophocytes that can mediate lysis of certain tumor cells and virue-infected cells without previous activation. They can also Natural killer (IMC) cells are lyophocytes and they cell self-time temperature of the light cell lectic-liker receptor family; also called NIG2 family; which is a group of transmembrane proteins preferentially expressed in NK cells. This family of proteins is characterized by the type in membrane orientation and the presence of a C-type lectin domain. This protein forms a complex with another family member; KIRD1: D94; and has been implicated in the recognition of the MHC class I HLAE-molecules in NK cells. The genes of NKG2 family members form a killer cell lectin-like receptor gene cluster on chromosome 12. Multiple alternatively spliced transcript variants encoding distinct isoforing disti

Usage Research use only

Unconjugated
All DIMA recombinant antibodies are genuinely generated by DIMA Biotech. They are all under patent application. Any protein sequencing or reverse engineering attempt is prohibited. We are actively scrutinizing all patent application to ensure no IP infringement. DIMA Disclaimer




Figure 1. Flow cytometry analysis with Anti-NKG2A (DMC298) on HEK293 cells transfected with human NKG2A (Blue histogram) or HEK293 transfected with irrelevant protein (Red histogram).

Address: Wuhan institute of Biotechnology B7, Biolake No.666 Gaoxin Road, Wuhan, Hubei, China Telephone: +1 2409940618(USA) /+86-18062749453(China) /+86-400-006-0995(China)

Email: info@dimabio.com Website: www.dimabio.com

