

PRODUCT INFORMATION

Warning: Undefined variable ShasAttributeValueDescription in C\u00e4wwroot\u00fcmirror.dimabio.com\u00e4wp-content\plugins\u00e4woocommerce-print-products\u00e4yublic\class-woocommerce-print-products-public.php on line 2806 |\u00e4\u

Conjugate Synonyms

Unconjugated
NKG2D;CD314;KLRK1;NK cell receptor D

Applications ELISA; Flow Cyt

Recommend Dilutions ELISA 1:5000-10000; Flow Cyt 1:100

Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution. Formulation & Reconstitution

Human IgG4 - Kappa IgG type Reactivity Target NKG2D Uniprot ID

Anti-NKG2D (tesnatilimab biosimilar) mAb Description

Delivery

In Stock
Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature. Storage & Sh

Background Research grade biosimilar. Not for use in therapeutic or diagnostic procedures for humans or animals

Usage

Anti-NKG2D (tesnatilimab biosimilar) mAb ELISA

0.2 µg of Human NKG2D, mFc Tagged protein per well





Figure 1. ELISA plate pre-coated by 2 μ g/ml (100 μ l/well) Human NKG2D, mFc tagged protein ([getskuurl sku="PME100079"]) can bind Anti-NKG2D Neutralizing antibody (BME100039) in a linear range of 0.64-400 ng/ml.



Email: info@dimabio.com Website: www.dimabio.com






Figure 2. Flow cytometry analysis with 15 μg/mL Anti-NKG2D (tesnatilimab) mAb (BME100039) on HEK293 cells transfected with Human NKG2D protein and Human DAP10 protein (Blue histogram) or HEK293 transfected with irrelevant protein (Red histogram).

Email: info@dimabio.com Website: www.dimabio.com

