

PRODUCT INFORMATION

| | |
|---|---|
| Common Name | Warning: Undefined variable \$hasAttributeValueDescription in C:\wwwroot\mirror\dimabio.com\wp-content\plugins\woocommerce-print-products\public\class-woocommerce-print-products-public.php on line 2806 PF-04518600 |
| Conjugate | Unconjugated |
| Synonyms | TNFRSF4;OX40;CD134;OX40L receptor;ACT35;TXGP1L |
| Applications | Flow Cyt |
| Recommended Dilutions | Flow Cyt 1:100 |
| Formulation & Reconstitution | Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution. |
| Host Species | Homo sapiens |
| IgG type | Human IgG2 - Kappa |
| Reactivity | Human |
| Target | OX40 |
| Uniprot ID | P43489 |
| Description | Anti-OX40(ivuxolimab biosimilar)mAb |
| Delivery | In Stock |
| Yefei Storage | Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature. |
| Background | Research grade biosimilar. Not for use in therapeutic or diagnostic procedures for humans or animals. |
| Usage | Research use only |




Figure 1. Flow cytometry analysis with Anti-OX40 mAb 1 μ g/ml (ivuxolimab) on HEK293 cells transfected with human OX40 (Blue histogram) or HEK293 transfected with irrelevant protein (Red histogram).

