

## **PRODUCT INFORMATION**

Warning: Undefined variable ShasAttributeValueDescription in C:\www.root\mirror.dimablo.com\wp-content\plugins\woocommerce-print-products\public\class-woocommerce-print-products-public.php on line 2806 LY-3462817,LY 3462817,LY 3462817

Conjugate

Unconjugated PD1, PDCD1, CD279, SLEB2, hPD-1, hPD-I, hSLE1 Synonyms

Applications ELISA

Recommend Dilutions

Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution. Formulation & Reconstitution

Human IgG1(E356D,M358L) - kappa IgG type

Reactivity Target PD-1 Uniprot ID

Description Anti-PD-1(peresolimab biosimilar) mAb

Delivery

Background


In Stock
Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized antibodies are shipped at ambient temperature.

Research grade biosimilar. Not for use in therapeutic or diagnostic procedures for humans or animals

Usage

## Anti-PD-1(peresolimab biosimilar) mAb ELISA

0.2 μg of Human PD-1, mFc-His tagged protein per well



Anti-PD-1(peresolimab biosimilar) mAb (ng/mL)

Figure 1. ELISA plate pre-coated by 2  $\mu$ g/mL (100  $\mu$ L/well) Human PD-1 Protein, mFc-His tag (PME100025) can bind Anti-PD-1(peresolimab biosimilar) mAb (BME100615) in a linear range of 3.2–80 ng/mL.

Email: info@dimabio.com Website: www.dimabio.com

Address: Wuhan institute of Biotechnology B7, Biolake No.666 Gaoxin Road, Wuhan, Hubei, China Telephone: +1 2409940618(USA) /+86-18062749453(China) /+86-400-006-0995(China)

