

**PRODUCT INFORMATION**

<b>Clone ID</b>	DM87
<b>Target</b>	PSCA
<b>Synonyms</b>	PSCA;UNQ206;PRO232
<b>Host Species</b>	Rabbit
<b>Description</b>	Anti-PSCA antibody(DM87); Rabbit mAb
<b>Delivery</b>	3~4 weeks
<b>Uniprot ID</b>	O43653
<b>IgG type</b>	Rabbit IgG
<b>Clonality</b>	Monoclonal
<b>Reactivity</b>	Human
<b>Applications</b>	ELISA; Flow Cyt; IHC
<b>Recommended Dilutions</b>	ELISA 1:5000-10000; Flow Cyt 1:100; IHC 1:100
<b>Purification</b>	Purified from cell culture supernatant by affinity chromatography
<b>Formulation &amp; Reconstitution</b>	Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.
<b>Storage &amp; Shipping</b>	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.
<b>Background</b>	This gene encodes a glycosylphosphatidylinositol-anchored cell membrane glycoprotein. In addition to being highly expressed in the prostate it is also expressed in the bladder; placenta; colon; kidney; and stomach. This gene is up-regulated in a large proportion of prostate cancers and is also detected in cancers of the bladder and pancreas. This gene includes a polymorphism that results in an upstream start codon in some individuals; this polymorphism is thought to be associated with a risk for certain gastric and bladder cancers. Alternative splicing results in multiple transcript variants.
<b>Usage</b>	Research use only
<b>Conjugate</b>	Unconjugated
<b>DIMA Disclaimer</b>	All DIMA recombinant antibodies are genuinely generated by DIMA Biotech. They are all under patent application. Any protein sequencing or reverse engineering attempt is prohibited. We are actively scrutinizing all patent application to ensure no IP infringement.






Figure 1. ELISA plate pre-coated by 2  $\mu\text{g/ml}$  (100  $\mu\text{l/well}$ ) Human PSCA protein, hFc tagged protein PME100084 can bind Rabbit anti-PSCA monoclonal antibody (clone: DM87) in a linear range of 1-100 ng/ml.




Figure 2. Expi 293 cell line transfected with irrelevant protein (A) and human PSCA (B) were surface stained with Rabbit anti-PSCA monoclonal antibody 1 $\mu\text{g/ml}$  (clone: DM87) followed by Alexa 488-conjugated anti-rabbit IgG secondary antibody.



**DME100087**




Figure 3. DME100087 at 10 $\mu$ g/ml staining PSCA in A431 xenografts in nude mice by IHC (SKU# DME100087)

DIMABIO CONFIDENTIAL

