

## **PRODUCT INFORMATION**

Warning . Undefined variable \$hasAttributeValueDescription in Calwawroottmirror.dimablo.com/wp-content\plugins/woocommerce-print-products-public.php on line 2806 bm174 Clone ID

ROR2;NTRKR2 Synonyme Host Species

Rabbit Anti-ROR2 antibody(DM174); Rabbit mAb

Description Delivery In Stock

Uniprot ID Q01974 lgG type Rabbit IgG Clonality Monoclonal Reactivity Human ELISA; Flow Cyt; WB Application

Recommend Dilutions ELISA 1:5000-10000; Flow Cyt 1:100; WB 1:1000

Purification Purified from cell culture supernatant by affinity chromatography

Formulation & Reconstitution Lyophilized from sterile PBS, pH 7.4. Normally 5 % – 8% trehalose is added as protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution.

Storage & Shipping

specific instructions of reconstitution.

Store at -20°C to .80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.

The protein encoded by this gene is a receptor protein byrosine kinase and type it transmembrane protein that belongs to the ROB subfamily of cell surface receptors. The protein may be involved in the early formation of the chondrocytes and may be required for cartilage and growth plate development. Mutations in this gene can cause brachydactyly type B; a seleted ild disorder characterized by hypoplasia applies of distall phalanges and nails. In addition; mutations in this gene can cause the autosomal recessive form of Robinow syndrome; which is characterized by skeletal dysplasia with generalized limb bone shortening; segmental defects of the spine; brachydactyly; and a dysmorphic facial appearance.

Research use only Background

Usage Coniugate

All DIMA recombinant antibodies are genuinely generated by DIMA Biotech. They are all under patent application. Any protein sequencing or reverse engineering attempt is prohibited. We are actively scrutinizing all patent application to ensure no IP infringement. DIMA Disclaime




Figure 1. ROR2 protein is highly expressed on the surface of HEK293 cell membrane. Flow cytometry analysis with Anti-ROR2 (DM174) on HEK293 cells transfected with human ROR2 (Blue histogram) or HEK293 transfected with irrelevant protein (Red histogram), and Isotype antibody on HEK293 transfected with irrelevant protein (Orange histogram).

Email: info@dimabio.com Website: www.dimabio.com

Address: Wuhan institute of Biotechnology B7, Biolake No.666 Gaoxin Road, Wuhan, Hubei, China Telephone: +1 2409940618(USA) /+86-18062749453(China) /+86-400-006-0995(China)








Figure 2. Flow cytometry analysis of antigen binding of rabbit anti-human ROR2 mAb(DME100174).

(A) DME100174 does not bind to Jurkat cells that do not express ROR2. (B) A clear peak shift of DME100174 was seen compared to the control when incubated with ROR2-expressing Raji cells, indicating strong binding of DME100174 to ROR2. Antibodies were incubated at 5  $\mu$ g/mL.



Figure 3.Anti-ROR2 antibody (SKU# DME100174) at 1/1000 dilution

Lane: RAJI(human Burkitt's lymphoma B lymphocyte), whole cell lysate

Secondary: Goat Anti-Rabbit IgG H&L (HRP) at 1/5000 dilution

Predicted band size: 105 kDa Observed band size: 140 kDa

> Email: info@dimabio.com Website: www.dimabio.com

